

RATE OF CONVERGENCE OF SPACE TIME APPROXIMATIONS FOR STOCHASTIC EVOLUTION EQUATIONS

ISTVÁN GYÖNGY AND ANNIE MILLET

ABSTRACT. Stochastic evolution equations in Banach spaces with unbounded nonlinear drift and diffusion operators driven by a finite dimensional Brownian motion are considered. Under some regularity condition assumed for the solution, the rates of convergence of various numerical approximations are estimated under strong monotonicity and Lipschitz conditions. The abstract setting involves general consistency conditions and is then applied to a class of quasilinear stochastic PDEs of parabolic type.

1. INTRODUCTION

Let $V \hookrightarrow H \hookrightarrow V^*$ be a *normal triple* of spaces with dense and continuous embeddings, where V is a separable and reflexive Banach space, H is a Hilbert space, identified with its dual by means of the inner product in H , and V^* is the dual of V .

Let $W = \{W(t) : t \geq 0\}$ be a d_1 -dimensional Brownian motion carried by a stochastic basis $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, P)$. Consider the stochastic evolution equation

$$u(t) = u_0 + \int_0^t A(s, u(s)) ds + \sum_{k=1}^{d_1} \int_0^t B_k(s, u(s)) dW^k(s), \quad t \in [0, T] \quad (1.1)$$

in the triple $V \hookrightarrow H \hookrightarrow V^*$, with a given H -valued \mathcal{F}_0 -measurable random variable u_0 , and given operators A and $B = (B_k)$, mapping $[0, \infty) \times \Omega \times V$ into V^* and $H^{d_1} := H \times \cdots \times H$, respectively. Let \mathcal{P} denote the σ -algebra of the predictable subsets of $[0, \infty) \times \Omega$, and let $\mathcal{B}(V)$, $\mathcal{B}(H)$ and $\mathcal{B}(V^*)$ be the Borel σ -algebras of V , H and V^* , respectively. Assume that A and B_k are $\mathcal{P} \otimes \mathcal{B}(V)$ -measurable with respect to the σ -algebras $\mathcal{B}(V^*)$ and $\mathcal{B}(H)$, respectively.

It is well-known that for any $T > 0$ equation (1.1) admits a unique solution u if A is *hemicontinuous* in $v \in V$, and (A, B) satisfies a *monotonicity*, *coercivity* and a *linear growth* condition (see [10], [13] and [16]). In [7] it is shown that under these conditions the solutions of various implicit and explicit schemes converge to u . In [8] the rate of convergence of implicit Euler approximations is estimated under more restrictive hypotheses: A and B satisfy a strong monotonicity condition, A is Lipschitz continuous in $v \in V$, and the solution u satisfies some regularity conditions.

1991 *Mathematics Subject Classification*. Primary: 60H15 Secondary: 65M60 .

Key words and phrases. Stochastic evolution equations, monotone operators, coercivity, space time approximations, Galerkin method, wavelets, finite elements.

This paper was written while the first named author was visiting the University of Paris 1. The research of this author is partially supported by EU Network HARP.

The research of the second named author is partially supported by the research project BMF2003-01345.

Then Theorem 3.4 from [8] in the case of time independent operators A and B reads as follows. For the implicit Euler approximation u^τ , corresponding to the mesh size $\tau = T/m$ of the partition of $[0, T]$, one has

$$E \max_{i \leq m} |u(i\tau) - u^\tau(i\tau)|_H^2 + \tau E \sum_{i \leq m} \|u(i\tau) - u^\tau(i\tau)\|_V^2 \leq C\tau^{2\nu},$$

where C is a constant, independent of τ , and $\nu \in]0, \frac{1}{2}]$ is a constant from the regularity condition imposed on u .

In this paper, we study space and space-time approximations schemes for equation (1.1) in a general framework. In order to obtain rate of convergence estimates we need to require more regularity from the solution u of equation (1.1) than what we can express in terms of the spaces V and H . Therefore in our setup we introduce additional Hilbert spaces \mathcal{V} and \mathcal{H} such that $\mathcal{V} \hookrightarrow \mathcal{H} \hookrightarrow V$, where \hookrightarrow denotes continuous embeddings. In examples these are Sobolev spaces such that \mathcal{H} and \mathcal{V} satisfy stronger differentiability conditions than V and \mathcal{H} , respectively. Our regularity conditions on the solution u are introduced in section 2 and labeled as **(R1)** and **(R2)**. In connection with these, we introduce also condition **(R3)**, requiring more regularity from A and B . Furthermore, condition **(R4)** on Hölder continuity in time of A and B is needed for schemes involving time discretization. We collect these conditions in Assumption 2.3 and call them *regularity conditions*.

In order to formulate ‘space discretizations’, we consider for any integer $n \geq 1$ a normal triple

$$V_n \hookrightarrow H_n \hookrightarrow V_n^*, \quad (1.2)$$

the ‘discrete’ counterpart of $V \hookrightarrow H \hookrightarrow V^*$, and a bounded linear operator

$$\Pi_n : V \rightarrow V_n,$$

connecting V to V_n . We have in mind discrete Sobolev spaces, wavelets and finite elements spaces, as examples for V_n .

The space discretization scheme for equation (1.1) is a stochastic evolutionary equation in the triple (1.2). We define it by replacing the operators A , B and the initial value u_0 in equation (1.1) by some $\mathcal{P} \otimes \mathcal{B}(V_n)$ -measurable operators

$$A^n : [0, \infty[\times \Omega \times V_n \rightarrow V_n^*, \quad B^n : [0, \infty[\times \Omega \times V_n \rightarrow H_n^{d_1}$$

and by an H_n -valued \mathcal{F}_0 -measurable random variable u_0^n , respectively, such that A^n and B^n satisfy in the triple (1.2) the strong monotonicity condition, the linear growth condition, A^n is hemicontinuous and B^n is Lipschitz continuous in $v \in V_n$. These are the conditions **(S1)**–**(S4)** in Assumption 3.1, which imply, in particular, the existence and uniqueness of a solution u^n to our scheme. We relate A_n and B_n to A and B via a *consistency condition*, **(Cn)** below. Then assuming **(S1)**–**(S4)**, under the regularity and consistency conditions **(R1)**, **(R3)** and **(Cn)** we have

$$E \sup_{0 \leq t \leq T} |\Pi_n u(t) - u^n(t)|_{H_n}^2 + E \int_0^T \|\Pi_n u(t) - u^n(t)\|_{V_n}^2 dt \leq CE |\Pi_n u_0 - u_0^n|_{H_n}^2 + C\varepsilon_n^2,$$

where C is a constant, independent of n , and $\varepsilon_n > 0$ is a constant from **(Cn)**. This is Theorem 3.1 below, our main result on the accuracy of approximations by space discretizations.

For an integer $m \geq 1$ we consider the grid $\{t_i = i\tau : 0 \leq i \leq m\}$ with mesh-size $\tau = T/m$. We define on this grid the space-time implicit and the space-time explicit approximations, $\{u_i^{n,\tau}\}_{i=0}^m$ and $\{u_{\tau,i}^n\}_{i=0}^m$, respectively, by

$$\begin{aligned} u_{i+1}^{n,\tau} &= u_i^{n,\tau} + \tau A_{i+1}^{n,\tau}(u_{i+1}^{n,\tau}) + \sum_k B_{k,i}^{n,\tau}(u_i^{n,\tau})(W^k(t_{i+1}) - W^k(t_i)), \\ u_{\tau,i+1}^n &= u_{\tau,i}^n + \tau A_i^{n,\tau}(u_{\tau,i}^n) + \sum_k B_{k,i}^{n,\tau}(u_{\tau,i}^n)(W^k(t_{i+1}) - W_k(t_i)), \end{aligned}$$

for $i = 0, \dots, m-1$ with some V_n -valued \mathcal{F}_0 -measurable random variables $u_0^{n,\tau}$ and $u_{0,\tau}^n$, and with some $\mathcal{F}_{t_i} \otimes \mathcal{B}(V)$ -measurable operators

$$A_i^{n,\tau} : \Omega \times V_n \rightarrow V_n^*, \quad B_{k,i}^{n,\tau} : \Omega \times V_n \rightarrow H_n^{d_1},$$

such that $A_i^{n,\tau}$, $B_{k,i}^{n,\tau}$ satisfy strong monotonicity and linear growth conditions and $A_i^{n,\tau}$ is Lipschitz continuous in $v \in V_n$. These conditions, listed as **(ST1)**–**(ST3)** in Assumption 4.1 below, are similar to conditions **(S1)**–**(S3)**, except that instead of the hemicontinuity, the much stronger assumption of Lipschitz continuity is assumed on $A_i^{n,\tau}$. The operators $A_i^{n,\tau}$ and $B_{k,i}^{n,\tau}$ are related to A and B by a consistency condition **(Cn τ)** stated below. Then if $\sup_{n,m} E|u_0^{n,\tau}|_{H_n}^2 < \infty$ and equation (1.1) satisfies the regularity conditions **(R1)**–**(R4)** from Assumption 2.3, we have the estimate

$$\begin{aligned} E \sup_{0 \leq i \leq m} |\Pi_n u(t_i) - u_i^{n,\tau}|_{H_n}^2 + E \sum_{0 \leq i \leq m} \|\Pi_n u(t_i) - u_i^{n,\tau}\|_{V_n}^2 \tau \\ \leq CE |\Pi_n u_0 - u_0^{n,\tau}|_{H_n}^2 + C(\tau^{2\nu} + \varepsilon_n^2), \end{aligned}$$

with a constant C , independent of n and τ , where $\nu \in]0, \frac{1}{2}]$ is the Hölder exponent from condition **(R4)** on the regularity of the operators A and B in time, and ε_n is from **(Cn τ)**. This is Theorem 4.4, our main result on implicit space-time approximations. In our main result, Theorem 5.2, on the explicit space-time approximations we have the same estimate for $u_{\tau,i}^n$ in place of $u_i^{n,\tau}$ if, in addition to the conditions of Theorem 4.4, as in [7], a *stability relation* between the time mesh τ and a space approximation parameter is satisfied.

Finally, we present as examples a class of quasi-linear stochastic partial differential equations (SPDEs) and linear SPDEs of parabolic type. We show that they satisfy the conditions of the abstract results, Theorems 3.1, 4.4 and 5.2, when we use wavelets, or finite differences. In particular, we obtain rate of convergence results for space and space-time approximations of linear parabolic SPDEs, among them for the Zakai equation of nonlinear filtering. We would like to mention that as far as we know, discrete Sobolev spaces are applied first in [18] to space discretizations and explicit space-time discretizations of linear SPDEs, and it inspired our approach to finite difference schemes. Our abstract results can also be applied to finite elements approximations. To keep down the size of the paper we will consider such applications elsewhere.

We denote by K, L, M and r some fixed constant, and by C some constants which, as usual, can change from line to line. For given constants $a \in \mathbb{R}^k$ the notation $C(a)$ means that the constant depends on a . Finally, when $(X, |\cdot|_X)$ and $(Y, |\cdot|_Y)$ denote two Banach spaces such that X is continuously embedded in Y , given $y \in Y$ the inequality $|y|_X < +\infty$ means that $y \in X$.

2. CONDITIONS ON EQUATION (1.1) AND ON THE APPROXIMATION SPACES

2.1. Conditions on equation (1.1). Let $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, P)$ be a stochastic basis, satisfying the usual conditions, i.e., $(\mathcal{F}_t)_{t \geq 0}$ is an increasing right-continuous family of sub- σ -algebras of \mathcal{F} such that \mathcal{F}_0 contains every P -null set. Let $W = \{W(t) : t \geq 0\}$ be a d_1 -dimensional Wiener martingale with respect to $(\mathcal{F}_t)_{t \geq 0}$, i.e., W is an \mathcal{F}_t -adapted Wiener process with values in \mathbb{R}^{d_1} such that $W(t) - W(s)$ is independent of \mathcal{F}_s for all $0 \leq s \leq t$. We use the notation \mathcal{P} for the sigma-algebra of predictable subsets of $[0, \infty) \times \Omega$. If V is a Banach space then $\mathcal{B}(V)$ denotes the sigma-algebra generated by the (closed) balls in V .

Let V be a separable reflexive Banach space embedded densely and continuously into a Hilbert space H , which is identified with its dual H^* by means of the inner product (\cdot, \cdot) in H . Thus we have a *normal triple*

$$V \hookrightarrow H \hookrightarrow V^*,$$

where $H \hookrightarrow V^*$ is the adjoint of the embedding $V \hookrightarrow H$. Thus $\langle v, h \rangle = (v, h)$ for all $v \in V$ and $h \in H^* = H$, where $\langle v, v^* \rangle = \langle v^*, v \rangle$ denotes the duality product of $v \in V$, $v^* \in V^*$, and (h_1, h_2) denotes the inner product of $h_1, h_2 \in H$. We assume, without loss of generality, that $|v|_H \leq \|v\|_H$ for all $v \in V$, where $|\cdot|_H$ and $\|\cdot\|_V$ denote the norms in H and V , respectively. For elements u from a normed space \mathbb{U} the notation $|u|_{\mathbb{U}}$ means the norm of u in \mathbb{U} .

Let A and $B = (B_k)_{k=1}^{d_1}$ be $\mathcal{P} \otimes \mathcal{B}(V)$ -measurable mappings from $[0, \infty) \times \Omega \times V$ into V^* and H^{d_1} , respectively. Given an H -valued \mathcal{F}_0 -measurable random variable u_0 consider the initial value problem

$$du(t) = A(t, u(t)) dt + \sum_k B_k(t, u(t)) dW^k(t), \quad u(0) = u_0 \quad (2.1)$$

on a fixed time interval $[0, T]$.

Assumption 2.1. *The operators A and B satisfy the following conditions.*

(i) (*Monotonicity of (A, B)*) *Almost surely for all $t \in [0, T]$ and $u, v \in V$,*

$$2\langle u - v, A(t, u) - A(t, v) \rangle + \sum_k |B_k(t, u) - B_k(t, v)|_H^2 \leq K|u - v|_H^2,$$

(ii) (*Coercivity of (A, B)*) *Almost surely for all $t \in [0, T]$ and $u, v \in V$,*

$$2\langle u, A(t, u) \rangle + \sum_k |B_k(t, u)|_H^2 + \mu\|u\|_V^2 \leq K|u|_H^2 + f(t), \quad (2.2)$$

(iii) (*Growth conditions on A and B*) *Almost surely for all $t \in [0, T]$ and $u \in V$,*

$$|A(t, u)|_{V^*}^2 \leq K_1\|u\|_V^2 + f(t), \quad \sum_k |B_k(t, u)|_H^2 \leq K_2\|u\|_V^2 + f(t),$$

(iv) (*Hemicontinuity of A*) *Almost surely for all $t \in [0, T]$ and $u, v, w \in V$,*

$$\lim_{\varepsilon \rightarrow 0} \langle w, A(t, u + \varepsilon v) \rangle = \langle w, A(t, u) \rangle, \quad (2.3)$$

where $\mu > 0$, $K \geq 0$, $K_1 \geq 0$ and $K_2 \geq 0$ are some constants, and f is a nonnegative (\mathcal{F}_t) -adapted stochastic process such that

$$E \int_0^T f(t) dt < \infty. \quad (2.4)$$

The following definition of solution is classical.

Definition 2.1. An H -valued adapted continuous process $u = \{u(t) : t \in [0, T]\}$ is a solution to equation (1.1) on $[0, T]$ if almost surely $u(t) \in V$ for almost every $t \in [0, T]$,

$$\int_0^T \|u(t)\|_V^2 dt < \infty,$$

and

$$(u(t), v) = (u_0, v) + \int_0^t \langle A(s, u(s)), v \rangle ds + \sum_k \int_0^t (B_k(s, u(s)), v) dW^k(s)$$

holds for all $t \in [0, T]$ and $v \in V$. We say that the solution to (2.1) on $[0, T]$ is unique if for any solutions u and v to (2.1) on $[0, T]$ we have

$$P\left(\sup_{t \in [0, T]} |u(t) - v(t)|_H > 0\right) = 0.$$

The following result is well known, see [10], [13], [16].

Theorem 2.2. Let Assumption 2.1 hold. Then (2.1) has a unique solution u . Moreover, if $E|u_0|_H^2 < \infty$, then

$$\begin{aligned} E \sup_{t \in [0, T]} |u(t)|_H^2 + E \int_0^T \|u(s)\|_V^2 ds \\ \leq CE|u_0|_H^2 + CE \int_0^T (f(t) + g(t)) dt < \infty, \end{aligned} \quad (2.5)$$

where C is a constant depending only on the constants λ , K and K_2 .

If Assumption 2.1 is satisfied then one can also show the convergence of approximations, obtained by various discretization schemes, to the solution u (see [7]). To estimate the rate of convergence of implicit time discretization schemes the following stronger assumption on A and B are used in [8]

Assumption 2.2. The operators A , B satisfy the following conditions almost surely.

(1) (Strong monotonicity) For all $t \in [0, T]$, $u, v \in V$,

$$\begin{aligned} 2\langle u - v, A(t, u) - A(t, v) \rangle + \sum_k |B_k(t, u) - B_k(t, v)|_H^2 \\ \leq -\lambda \|u - v\|_V^2 + L|u - v|_H^2, \end{aligned}$$

(2) (Growth conditions on A and B) For all $t \in [0, T]$, $u \in V$,

$$|A(t, u)|_{V^*}^2 \leq K_1 \|u\|_V^2 + f(t), \quad \sum_k |B_k(t, u)|_H^2 \leq K_2 \|u\|_V^2 + g(t). \quad (2.6)$$

(3) (Lipschitz condition on A) For all $t \in [0, T]$, $u, v \in V$,

$$|A(t, u) - A(t, v)|_{V^*}^2 \leq L_1 \|u - v\|_V^2, \quad (2.7)$$

where $\lambda > 0$, $K \geq 0$, $K_1 \geq 0$, $K_2 \geq 0$ are constants, and f and g are non-negative adapted processes satisfying (2.4)

Remark 2.3. *It is easy to see that due to (1)–(2), the coercivity condition (2.2) holds with $\mu = \lambda/2$ and a constant $K = K(\lambda, L, K_2)$.*

Remark 2.4. *It is easy to show that (1) and (3) imply that $B = (B_k)$ is Lipschitz continuous in $u \in V$, i.e., almost surely*

$$\sum_k |B_k(t, u) - B_k(t, v)|_H^2 \leq L_2 \|u - v\|_V^2 \quad \text{for all } u, v \in V, t \in [0, T] \quad (2.8)$$

where L_2 is a constant depending on λ , L and L_1 .

In order to prove rate of convergence estimates for the approximation schemes presented in this paper, we need to impose additional *regularity conditions* on equation (2.1) and on the solution u . Therefore we assume that there exist some separable Hilbert spaces \mathcal{V} and \mathcal{H} such that

$$\mathcal{V} \hookrightarrow \mathcal{H} \hookrightarrow V,$$

where \hookrightarrow means continuous embedding, and introduce the following conditions.

Let K, M denote some constants, fixed throughout the paper.

Assumption 2.3. (*Regularity conditions*)

(R1) *There is a unique solution u of (2.1), it takes values in \mathcal{V} for $dt \times P$ -almost every $(t, \omega) \in [0, T] \times \Omega$, $u_0 \in V$ and*

$$E \|u_0\|_V^2 < \infty, \quad E \int_0^T |u(t)|_V^2 dt =: r_1 < \infty. \quad (2.9)$$

(R2) *There is a unique solution u of (2.1), it has an \mathcal{H} -valued stochastic modification, denoted also by u , such that*

$$\sup_{t \in [0, T]} E |u(t)|_{\mathcal{H}}^2 =: r_2 < \infty.$$

(R3) *Almost surely $A(t, v) \in V$, $B_k(t, u) \in V$ and*

$$\|A(t, v)\|_V^2 \leq K |v|_V^2 + \xi(t), \quad \sum_k \|B_k(t, u)\|_V^2 \leq K |u|_{\mathcal{H}}^2 + \eta(t) \quad (2.10)$$

for all $t \in [0, T]$ $v \in \mathcal{V}$ and $u \in \mathcal{H}$, where ξ and η are non-negative processes such that for some constant M

$$E \int_0^T \xi(t) dt \leq M, \quad \sup_{t \in [0, T]} E \eta(t) \leq M.$$

(R4) (*Time regularity of A, B*) *There exists a constant $\nu \in]0, \frac{1}{2}]$ and a non-negative random variable η such that $E \eta \leq M$, and almost surely*

$$(i) \quad \|A(s, v) - A(t, v)\|_V^2 \leq (K |v|_V^2 + \eta) |t - s|^{2\nu} \quad \text{for } v \in \mathcal{V}, \quad (2.11)$$

$$(ii) \quad \sum_k |B_k(s, u) - B_k(t, u)|_V^2 \leq (K |u|_V^2 + \eta) |t - s|^{2\nu} \quad \text{for } u \in \mathcal{V},$$

for all $0 \leq s < t \leq T$.

Remark 2.5. Assume conditions **(R1)** and **(R3)** from Assumption 2.3. Then the following statements hold.

(i) u has a V -valued continuous stochastic modification, denoted also by u , such that

$$E \sup_{t \in [0, T]} \|u(t)\|^2 \leq 3E\|u_0\|_V^2 + C(r_1 + M);$$

(ii) If condition **(R2)** from Assumption 2.3 also holds, then for $s, t \in [0, T]$,

$$E\|u(t) - u(s)\|_V^2 \leq C|t - s|(r_1 + r_2 + M), \quad (2.12)$$

where C is a constant depending only on T and on the constant K from (2.10).

Proof. Define

$$F(t) = \int_0^t A(s, u(s)) ds \quad \text{and} \quad G(t) = \sum_k \int_0^t B_k(s, u(s)) dW^k(s).$$

Notice that

$$\begin{aligned} E \int_0^T \|A(s, u(s))\|^2 ds &\leq KE \int_0^T |u(s)|_V^2 ds + E \int_0^T \xi(s) ds =: M_1 < \infty, \\ \sum_k \int_0^T E \|B_k(s, u(s))\|_V^2 ds &\leq KE \int_0^T |u(s)|_{\mathcal{H}}^2 ds + E \int_0^T \eta(s) ds =: M_2 < \infty. \end{aligned}$$

Hence F and G are V -valued continuous processes, and by Jensen's and Doob's inequalities

$$E \sup_{t \leq T} \|F(t)\|_V^2 \leq TM_1, \quad E \sup_{t \leq T} \|G(t)\|_V^2 \leq 4 \sum_k E \int_0^T \|B_k(s, u(s))\|_V^2 ds \leq 4M_2.$$

Consequently, the process $u_0 + F(t) + G(t)$ is a V -valued continuous modification of u , and statement (i) holds. Moreover, if **(R2)** also holds, then

$$\sup_{t \in [0, T]} \sum_k E \|B_k(s, u(s))\|_V^2 \leq K \sup_{t \in [0, T]} E |u(t)|_{\mathcal{H}}^2 + \sup_{t \in [0, T]} E \eta(t) := M_3 < +\infty,$$

and

$$\begin{aligned} E\|F(t) - F(s)\|_V^2 &\leq |t - s|M_1, \\ E\|G(t) - G(s)\|_V^2 &= \sum_k \int_s^t E \|B_k(r, u(r))\|_V^2 dr \leq |t - s|M_3 \end{aligned}$$

for any $0 \leq s \leq t \leq T$, which proves (ii). \square

2.2. Approximation spaces and operators Π_n . Let $V_n \hookrightarrow H_n \hookrightarrow V_n^*$ be a normal triple and $\Pi_n : V \rightarrow V_n$ be a bounded linear operator for each integer $n \geq 0$ such that for all $v \in H$ and $n \geq 0$

$$\|\Pi_n v\|_{V_n} \leq p \|v\|_V \quad (2.13)$$

with some constant p independent of $v \in V$ and n . Note that we do not require that the maps Π_n be orthogonal projections on the Hilbert space H .

We denote by $\langle v, w \rangle_n$ the duality between $v \in V_n$ and $w \in V_n^*$ and similarly by $(h, k)_n$ the inner product of $h, k \in H_n$. To lighten the notation, let $\|v\| := \|v\|_V$ denote the norm of v in V , $\|v\|_n := \|v\|_{V_n}$ the norm of v in V_n , $|u| := |u|_H$ the

norm of u in H , $|u|_n = |u|_{H_n}$ the norm of u in H_n , and finally $|w|_* := |w|_{V^*}$ and $|y|_{n^*} := |y|_{V_n^*}$ the norm of $w \in V^*$ in V^* and the norm of y in V_n^* , respectively.

For $r \geq 0$ let $H^r = W_2^r(\mathbb{R}^d)$ denote the closure of $C_0^\infty(\mathbb{R}^d)$ in the norm defined by

$$|\varphi|_{H^r}^2 = \sum_{|\gamma| \leq r} \int_{\mathbb{R}^d} |D^\gamma \varphi(x)|^2 dx.$$

In particular, $H^0 = L_2(\mathbb{R}^d)$.

The following basic examples will be used in the sequel. It describes spaces V_n, H_n and V_n^* and operators Π_n such that condition (2.13) is satisfied.

Example 2.6. Wavelet approximation. Let $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ be an orthonormal scaling function, i.e., a real-valued, compactly supported function, such that:

(i) there exists a sequence $(h_k)_{k \in \mathbb{Z}} \in l^2(\mathbb{Z})$ for which $\varphi(x) = \sum_k h_k \varphi(2x - k)$ in $L^2(\mathbb{R})$,

(ii) $\int \varphi(x - k) \varphi(x - l) dx = \delta_{k,l}$ for any $k, l \in \mathbb{Z}$.

We assume that the scaling function φ belongs to the Sobolev space $H^s(\mathbb{R}) := W_2^s(\mathbb{R})$ for sufficiently large integer $s > 0$.

For $d > 1$, $x = (x_1, \dots, x_d) \in \mathbb{R}^d$, set $\phi(x) = \varphi(x_1) \cdots \varphi(x_d)$ and for $j \geq 0$ and $k \in \mathbb{Z}^d$, set $\phi_{j,k}(x) = 2^{\frac{jd}{2}} \phi(2^j x - k) \in H^s := W_2^s(\mathbb{R}^d)$. For any integer $j \geq 0$, let H_j denote the closure in $L^2(\mathbb{R}^d)$ of the vector space generated by $(\phi_{j,k}, k \in \mathbb{Z}^d)$ and define the operator Π_j by

$$\Pi_j f = \sum_{k \in \mathbb{Z}^d} (f, \phi_{j,k}) \phi_{j,k}, \quad f \in L^2(\mathbb{R}^d),$$

$(,)$ denotes the scalar product in $L^2(\mathbb{R}^d)$.

Thus we have a sequence $H_j \subset H_{j+1}$ of closed subspaces of $L^2(\mathbb{R}^d)$ and orthogonal projections $\Pi_j : L^2(\mathbb{R}^d) \rightarrow H_j$ for $j \geq 0$. Assume, moreover that $\cup_{j=0}^\infty H_j$ is dense in $L^2(\mathbb{R}^d)$ and that φ is sufficiently regular, such that the inequalities

$$(Direct) \quad \|f - \Pi_j f\|_{H^r} \leq C 2^{-j(s-r)} \|f\|_{H^s}, \quad \forall f \in H^s, \quad (2.14)$$

$$(Converse) \quad \|\Pi_j f\|_{H^s} \leq C 2^{j(s-r)} \|f\|_{H^r}, \quad \forall f \in H^r \quad (2.15)$$

holds for fixed integers $0 \leq r \leq s$. The proof of these inequalities and more information on wavelets can be found, e.g., in [2].

Fix $r > 0$, set $H := L^2(\mathbb{R}^d)$, $V := H^r = W_2^r(\mathbb{R}^d)$, and identify H with its dual H^* by the help of the inner product in H . Then $V \hookrightarrow H^* \hookrightarrow V^*$ is a normal triple, where $H \equiv H^* \hookrightarrow V^*$ is the adjoint of the embedding $V \hookrightarrow H$. We define V_n as the normed space we get by taking the H^r norm on H_n . Since the H^r and H^0 norms are equivalent on H_n , the space V_n is complete, and obviously $V_n \hookrightarrow H_n \equiv H_n^* \hookrightarrow V_n^*$ is a normal triple, where H_n is identified with H_n^* via the inner product $(,)_n = (,)$ in H_n . Note that due to (2.15) we have (2.13) assuming that φ is sufficiently smooth.

Example 2.7. Finite differences – Discrete Sobolev spaces. Consider for fixed $h \in (0, 1)$ the grid

$$\mathbb{G} = h\mathbb{Z}^d = \{(k_1 h, k_2 h, \dots, k_d h) : k = (k_1, k_2, \dots, k_d) \in \mathbb{Z}^d\},$$

where \mathbb{Z} denotes the set of integers. Use the notation $\{e_1, e_2, \dots, e_d\}$ for the standard basis in \mathbb{R}^d . For any integer $m \geq 0$, let $W_{h,2}^m$ be the set of real valued functions v on \mathbb{G} with

$$|v|_{h,m}^2 := \sum_{|\alpha| \leq m} \sum_{z \in \mathbb{G}} |\delta_+^\alpha v(z)|^2 h^d < \infty,$$

where $\delta_{\pm i}^0$ is the identity and $\delta_{\pm}^\alpha = \delta_{\pm 1}^{\alpha_1} \delta_{\pm 2}^{\alpha_2} \dots \delta_{\pm d}^{\alpha_d}$ for multi-indices $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_d) \in \{0, 1, 2, \dots\}^d$ of length $|\alpha| := \alpha_1 + \dots + \alpha_d \geq 1$ is defined for by

$$\delta_{\pm i} v(z) := \pm \frac{1}{h} (v(z \pm h e_i) - v(z)).$$

We write also δ^α and δ_i in place of δ_+^α and δ_{+i} , respectively. Then $W_{h,2}^m$ with the norm $|\cdot|_{h,m}$ is a separable Hilbert space. It is the discrete counterpart of the Sobolev space $W_2^m(\mathbb{R}^d)$. Set $W_{h,2}^{-1} = (W_{h,2}^1)^*$, the adjoint of $W_{h,2}^1$, with its norm denoted by $|\cdot|_{h,-1}$. It is easy to see that $W_{h,2}^m \hookrightarrow W_{h,2}^{m-1}$ is a dense and continuous embedding,

$$\begin{aligned} |v|_{h,m-1} &\leq |v|_{h,m}, \\ |v|_{h,m} &\leq \frac{\kappa}{h} |v|_{h,m-1}, \end{aligned} \quad (2.16)$$

for all $v \in W_{h,2}^m$, $m \geq 0$ and $h \in (0, 1)$, where κ is a constant depending only on d . Notice that for $m \geq 1$

$$\langle v, u \rangle := \sum_{|\alpha| \leq m} \sum_{z \in \mathbb{G}} \delta^\alpha v \delta^\alpha u \leq C |v|_{h,m-1} |u|_{h,m+1} \quad \text{for all } v, u \in W_{h,2}^{m+1}$$

extends to a duality product between $W_{h,2}^{m-1}$ and $W_{h,2}^{m+1}$, which makes it possible to identify $W_{h,2}^{m-1}$ with $(W_{h,2}^{m+1})^*$.

Assume that $m > \frac{d}{2}$. Then by Sobolev's theorem on embedding $W_2^m := W_2^m(\mathbb{R}^d)$ into $\mathcal{C}(\mathbb{R}^d)$, there is a bounded linear operator $I : W_2^m(\mathbb{R}^d) \rightarrow \mathcal{C}(\mathbb{R}^d)$, such that $Iu = u$ almost everywhere on \mathbb{R}^d . Thus, identifying u with Iu , we can define the operator $R_h : W_2^m(\mathbb{R}^d) \rightarrow W_{h,2}^m$ by restricting the functions $u \in W_2^m$ onto $\mathbb{G} \subset \mathbb{R}^d$. Moreover, due to Sobolev's theorem,

$$\sum_{z \in \mathbb{G}} \sup_{x \in \mathcal{I}(z)} |u(x)|^2 h^d \leq p^2 |u|_{W_2^m}^2,$$

where $\mathcal{I}(z) := \{x \in \mathbb{R}^d : z_k \leq x_k \leq z_k + h, \quad k = 1, 2, \dots, d\}$ and p is a constant depending only on m and d . Hence obviously

$$|R_h u|_{h,0}^2 \leq p |u|_{W_2^m}^2 \quad \text{for all } u \in W_2^m. \quad (2.17)$$

Moreover, for every integer $l \geq 0$

$$|R_h u|_{h,l} \leq p |u|_{W_2^{m+l}} \quad \text{for all } u \in W_2^{m+l}, \quad (2.18)$$

with a constant p depending only on m , l and d . Thus setting

$$\begin{aligned} V_n &:= W_{h_n,2}^{m+l}, & H_n &:= W_{h_n,2}^{m+l-1}, & V_n^* &\equiv W_{h_n,2}^{m+l-2}, \\ \Pi_n &:= R_{h_n} \end{aligned}$$

for any sequence $\{h_n\}_{n=0}^\infty \subset (0, 1)$ and any integers $m > \frac{d}{2}$, $l \geq 0$ we get examples of approximation spaces.

When approximating differential operators by finite differences we need to estimate $D_i u - \delta_{\pm i} u$ in discrete Sobolev norms. For $d = 1$ we can estimate this as

follows. Let $l \geq 0$ be an integer and set $z_k := kh$ for $k \in \mathbb{Z}$. By the mean value theorem there exist z'_k and z''_k in $[z_k, z_k + lh]$ such that $\delta^l Du(z_k) = D^{l+1}u(z'_k)$ and $\delta^l \delta u(z_k) = D^{l+1}u(z''_k)$, where $D := \frac{d}{dx}$. Hence

$$\begin{aligned} |\delta^l(Du(z_k) - \delta u(z_k))|^2 &= |D^{l+1}u(z'_k) - D^{l+1}u(z''_k)|^2 = \left| \int_{z'_k}^{z''_k} D^{l+2}u(y) dy \right|^2 \\ &\leq lh \int_{z_k}^{z_k+lh} |D^{l+2}u(y)|^2 dy \end{aligned}$$

for $u \in C_0^\infty(\mathbb{R})$. Consequently,

$$|Du - \delta_\pm^l u|_{h,l} \leq lh |u|_{W_2^{l+2}(\mathbb{R})} \quad (2.19)$$

for $u \in C_0^\infty(\mathbb{R})$, and hence for all $u \in W_2^{l+2}(\mathbb{R})$. For $d > 1$ by similar calculation combined with Sobolev's embedding, we get that for $m > l + 2 + \frac{d-1}{2}$

$$|D_i u - \delta_{\pm i} u|_{h,l} \leq Ch |u|_{W_2^m} \quad (2.20)$$

for all $u \in W_2^m$, $h \in (0, 1)$, where C is a constant depending on l, m and d .

3. SPACE DISCRETIZATION

3.1. Description of the scheme. Consider for each integer $n \geq 1$ the problem

$$du^n(t) = A^n(t, u^n(t)) dt + \sum_k B^n(t, u^n(t)) dW^k(t), \quad u^n(0) = u_0^n, \quad (3.1)$$

in a normal triple $V_n \hookrightarrow H_n \hookrightarrow V_n^*$, satisfying the conditions of section 2.2, where u_0^n is an H_n -valued \mathcal{F}_0 -measurable random variable, and A^n and $B^n = (B_k^n)$ are $\mathcal{P} \otimes \mathcal{B}(V_n)$ -measurable mappings from $[0, \infty) \times \Omega \times V_n$ into V_n^* and $H_n^{d_1}$, respectively.

Assumption 3.1. *The operators A^n and B^n satisfy the following conditions.*

(S1) *(Strong monotonicity) There exist constants $\lambda > 0$ and L such that for all $n \geq 1$ almost surely*

$$\begin{aligned} 2\langle u - v, A^n(t, u) - A^n(t, v) \rangle_n + \sum_k |B_k^n(t, u) - B_k^n(t, v)|_{H_n}^2 + \lambda \|u - v\|_{V_n}^2 \\ \leq L \|u - v\|_{H_n}^2 \quad \text{for all } t \in [0, T], u, v \in V_n. \end{aligned}$$

(S2) *(Growth condition) Almost surely*

$$|A^n(t, v)|_{V_n^*}^2 \leq K_1 \|v\|_{V_n}^2 + f^n(t), \quad |B^n(t, v)|_{H_n}^2 \leq K_2 \|v\|_{V_n}^2 + g^n(t)$$

for all $t \in [0, T]$, $v \in V_n$ and $n \geq 1$, where K_1, K_2 are constants, independent of n , and f^n and g^n are non-negative stochastic processes such that

$$\sup_n E \int_0^T f^n(t) dt =: M_1 < \infty, \quad \sup_n E \int_0^T g^n(t) dt =: M_2 < \infty.$$

(S3) *(Hemicontinuity of A^n) For every $n \geq 1$, the operators A^n are hemicontinuous in $v \in V_n$, i.e., almost surely*

$$\lim_{\varepsilon \rightarrow 0} \langle A^n(t, v + \varepsilon u), w \rangle_n = \langle A^n(t, v), w \rangle_n$$

for all $t \in [0, T]$, $v, u, w \in V_n$.

(S4) (*Lipschitz condition on B^n*) Almost surely

$$\sum_k |B_k^n(t, u) - B_k^n(t, v)|_{H_n}^2 \leq L_B \|u - v\|_{V_n}^2$$

for all $t \in [0, T]$ and $u, v \in V_n$.

The solution to (3.1) is understood in the sense of Definition 2.1. Notice that **(S1)** - **(S2)** imply the coercivity condition

$$2\langle v, A^n(t, v) \rangle_n + \sum_k |B_k^n(t, v)|_{H_n}^2 + \frac{\lambda}{2} \|v\|_{V_n}^2 \leq C (|v|_{H_n}^2 + f^n(t) + g^n(t))$$

with a constant C depending on λ , L and K_2 .

Thus by Theorem 2.2 the conditions **(S1)**–**(S3)** ensure the existence of a unique solution u^n to (3.1), and if

$$\sup_n E|u_0^n|_{H_n}^2 < \infty, \quad (3.2)$$

then

$$\begin{aligned} E \sup_{0 \leq t \leq T} |u^n(t)|_{H_n}^2 + E \int_0^T \|u^n(t)\|_{V_n}^2 dt \\ \leq C \sup_n \left(E|u_0^n|_{H_n}^2 + E \int_0^T (f^n(t) + g^n(t)) dt \right) < \infty, \end{aligned} \quad (3.3)$$

where C is a constant depending only on λ , L and K_2 .

3.2. Rate of convergence of the scheme. We want to approximate $\Pi_n u$ by u^n . In order to estimate the accuracy of this approximation we need to relate the operators A and B to A^n and B^n , respectively. Therefore we assume the regularity condition **(R3)** from Assumption 2.3 and make the following *consistency* assumption.

Condition (Cn) (*Consistency*) There exist a sequence $(\varepsilon_n)_{n \geq 1}$ of positive numbers and a sequence $(\xi^n)_{n \geq 1}$ of non-negative adapted processes such that

$$\sup_n E \int_0^T \xi^n(t) dt \leq M < +\infty,$$

and almost surely $(t, \omega) \in [0, T] \times \Omega$

$$\begin{aligned} |\Pi_n A(t, v) - A^n(t, \Pi_n v)|_{V_n^*}^2 + \sum_k |\Pi_n B_k(t, v) - B_k^n(t, \Pi_n v)|_{H_n}^2 \\ \leq \varepsilon_n^2 (|v|_{V_n}^2 + \xi^n(t)) \end{aligned}$$

for all $t \in [0, T]$ and $v \in \mathcal{V}$.

Theorem 3.1. *Let Assumption 3.1, the regularity conditions **(R1)** and **(R3)** from Assumption 2.3, and the consistency condition **(Cn)** hold. Assume furthermore $\sup_n E|u_0^n|_{H_n}^2 < +\infty$. Then for $e^n(t) := \Pi_n u(t) - u^n(t)$,*

$$E \sup_{0 \leq t \leq T} |e^n(t)|_{H_n}^2 + E \int_0^T \|e^n(t)\|_{V_n}^2 dt \leq C_1 E|e^n(0)|_{H_n}^2 + C_2(r_1 + M)\varepsilon_n^2 \quad (3.4)$$

holds for all $n \geq 1$, where $C_1 = C_1(\lambda, L, T)$ and $C_2 = C_2(\lambda, L, L_B, T)$ are constants.

Proof. From equation (1.1) we deduce that for every $n \geq 1$,

$$\Pi_n u(t) = \Pi_n u_0 + \int_0^t \Pi_n A(s, u(s)) ds + \sum_k \int_0^t \Pi_n B_k(s, u(s)) dW^k(s).$$

Using Itô's formula

$$|e^n(t)|_n^2 = |e^n(0)|_n^2 + \sum_{i \leq 3} I_i(t), \quad (3.5)$$

where

$$\begin{aligned} I_1(t) &= 2 \int_0^t \langle e^n(s), \Pi_n A(s, u(s)) - A^n(s, u^n(s)) \rangle_n ds, \\ I_2(t) &= 2 \sum_k \int_0^t (e^n(s), \Pi_n B_k(s, u(s)) - B_k^n(s, u^n(s)))_n dW^k(s), \\ I_3(t) &= \sum_k \int_0^t |\Pi_n B_k(s, u(s)) - B_k^n(s, u^n(s))|_n^2 ds. \end{aligned}$$

We first prove

$$\sup_{0 \leq t \leq T} E|e^n(t)|_n^2 + E \int_0^T \|e^n(t)\|_n^2 dt \leq C_1 E|e^n(0)|_n^2 + C_2(r_1 + M)\varepsilon_n^2, \quad (3.6)$$

where $C_1 = C_1(\lambda, L, T)$ and $C_2 = C_2(\lambda, L, L_B, T)$ are constants. The strong monotonicity condition **(S1)** from Assumption 3.1 implies

$$I_1(t) + I_3(t) \leq -\lambda \int_0^t \|e^n(s)\|_n^2 ds + L \int_0^t |e^n(s)|_n^2 ds + \sum_{i=1,2} R_i(t), \quad (3.7)$$

where

$$\begin{aligned} R_1(t) &= \int_0^t 2 \langle e^n(s), \Pi_n A(s, u(s)) - A^n(s, \Pi_n u(s)) \rangle_n ds, \\ R_2(t) &= \sum_k \int_0^t \left[|\Pi_n B_k(s, u(s)) - B_k^n(s, u^n(s))|_n^2 \right. \\ &\quad \left. - |B_k^n(s, \Pi_n u(s)) - B_k^n(s, u^n(s))|_n^2 \right] ds. \end{aligned}$$

Schwarz's inequality and the consistency condition **(Cn)** imply that for every $n \geq 1$ and $t \in [0, T]$,

$$\begin{aligned} |R_1(t)| &\leq \frac{\lambda}{3} \int_0^t \|e^n(s)\|_n^2 ds + \frac{3}{\lambda} \int_0^t |\Pi_n A(s, u(s)) - A^n(s, \Pi_n u(s))|_{n^*}^2 ds \\ &\leq \frac{\lambda}{3} \int_0^t \|e^n(s)\|_n^2 ds + \frac{3}{\lambda} \varepsilon_n^2 \int_0^t (|u(s)|_V^2 + \xi^n(s)) ds. \end{aligned} \quad (3.8)$$

Schwarz's inequality, the consistency condition **(Cn)**, and the Lipschitz condition **(S4)** from Assumption 3.1 yield that for every $\alpha > 0$,

$$|R_2(t)| = \sum_k \int_0^t \left[|\Pi_n B_k(s, u(s)) - B_k^n(s, \Pi_n u(s))|_n^2 \right.$$

$$\begin{aligned}
 & + 2 \left(\Pi_n B_k(s, u(s)) - B_k^n(s, \Pi_n u(s)), B_k^n(s, \Pi_n u(s)) - B_k^n(s, u^n(s)) \right)_n \Big] ds \\
 & \leq (1 + \frac{1}{\alpha}) \int_0^t \sum_k |\Pi_n B_k(s, u(s)) - B_k^n(s, \Pi_n u(s))|_n^2 ds \\
 & \quad + \alpha \int_0^t \sum_k |B_k^n(s, \Pi_n u(s)) - B_k^n(s, u^n(s))|_n^2 ds \\
 & \leq (1 + \frac{1}{\alpha}) \varepsilon_n^2 \int_0^t (|u(s)|_V^2 + \xi^n(s)) ds + \alpha L_B \int_0^t \|e^n(s)\|_n^2 ds. \tag{3.9}
 \end{aligned}$$

Thus, for $\alpha L_B \leq \frac{\lambda}{3}$, taking expectations in (3.5) and (3.7)-(3.9) and using **(S1)** again, we deduce that

$$E|e^n(t)|_n^2 + \frac{\lambda}{3} E \int_0^t \|e^n(s)\|_n^2 ds \leq LE \int_0^t |e^n(s)|_n^2 ds + E|e^n(0)|_n^2 + C(r_1 + M)\varepsilon_n^2,$$

where $C = C(\lambda, L_B)$ is a constant. Since by (2.5) and (3.3)

$$\sup_{0 \leq t \leq T} E|e^n(t)|_n^2 < +\infty,$$

Gronwall's lemma gives

$$\sup_{0 \leq t \leq T} E|e^n(t)|_n^2 \leq e^{LT} (C(r_1 + M)\varepsilon_n^2 + E|e^n(0)|_n^2),$$

which in turn yields (3.6). We now prove (3.4). From (3.6)–(3.9) we deduce

$$\begin{aligned}
 E \sup_{0 \leq t \leq T} (I_1(t) + I_3(t)) & \leq LE \int_0^T |e^n(s)|_n^2 ds + \frac{2\lambda}{3} E \int_0^T \|e^n(s)\|_n^2 ds + C_2(r_1 + M) \varepsilon_n^2 \\
 & \leq C_1 E|e^n(0)|_n^2 + C_2(r_1 + M)\varepsilon_n^2. \tag{3.10}
 \end{aligned}$$

(Notice that by taking the supremum in both sides of (3.7) we cannot make use of the term with coefficient $-\lambda$ in the right-hand side of (3.7). This is why $2\lambda/3$ appears here as the sum of $\lambda/3$ from (3.8) and $\alpha L_B \leq \lambda/3$ from (3.9).) By Davies' inequality, (2.5), the Lipschitz condition **(S4)** on B^n , the consistency condition **(Cn)** and by the strong monotonicity condition **(S1)**,

$$\begin{aligned}
 E \sup_{0 \leq t \leq T} |I_2(t)| & \leq 6E \left(\int_0^T \sum_k \left| \left(e^n, \Pi_n B_k(u) - B_k^n(u^n) \right)_n \right|^2 ds \right)^{\frac{1}{2}} \\
 & \leq 6E \left\{ \sup_{0 \leq t \leq T} |e^n(t)|_n \left(\int_0^T \sum_k |\Pi_n B_k(u) - B_k^n(u^n)|_n^2 ds \right)^{\frac{1}{2}} \right\} \\
 & \leq \frac{1}{2} E \sup_{0 \leq t \leq T} |e^n(t)|_n^2 \\
 & \quad + 36E \sum_k \int_0^T \left[|\Pi_n B_k(u) - B_k^n(\Pi_n u)|_n^2 + |B_k^n(\Pi_n u) - B_k^n(u^n)|_n^2 \right] ds \\
 & \leq \frac{1}{2} E \sup_{0 \leq t \leq T} |e^n(t)|_n^2 + 36L_B E \int_0^T \|e^n(s)\|_n^2 ds + 36(r_1 + M)\varepsilon_n^2, \tag{3.11}
 \end{aligned}$$

where the argument s is omitted from most integrands. Thus, relations (3.5), (3.10), (3.11) and (3.6) yield

$$\frac{1}{2}E \sup_{0 \leq t \leq T} |e^n(t)|_n^2 \leq C_1 E |e^n(0)|_n^2 + C_2(r_1 + M)\varepsilon_n^2,$$

with some constants $C_1 = C_1(L, T)$ and $C_2 = C_2(\lambda, L, L_B, T)$, which completes the proof of (3.4). \square

3.3. Example. Consider the normal triples

$$V \hookrightarrow H^* \hookrightarrow V^*, \quad V_n \hookrightarrow H_n^* \hookrightarrow V_n^*$$

with the orthogonal projection $\Pi_n : H = L^2(\mathbb{R}^d) \rightarrow H_n$ from Example 2.6, where $V = W_2^r(\mathbb{R}^d)$ with $r > 0$. Set $\mathcal{H} = W_2^{r+\rho}(\mathbb{R}^d)$ and $\mathcal{V} = W_2^{r+l}(\mathbb{R}^d)$ for some $l > \rho \geq 0$.

Let A and $B = (B_k)$ be $\mathcal{P} \otimes \mathcal{B}(V)$ -measurable mappings from $[0, \infty[\times\Omega \times V$ into V^* and H^{d_1} , respectively, satisfying Assumptions 2.2 and 2.3. For $(t, \omega) \in [0, T] \times \Omega$ let $A^n(t, \omega, \cdot) : V^n \rightarrow V_n^*$ and $B^n(t, \omega, \cdot) : V^n \rightarrow H_n^{d_1}$ be defined by

$$\langle A^n(t, u, \omega), v \rangle_n = \langle A(t, u, \omega), v \rangle \quad \text{and} \quad B_k^n(t, \omega, u) = \Pi_n B_k(t, \omega, u) \quad (3.12)$$

for all $u, v \in V_n$, where $\langle \cdot, \cdot \rangle_n$ denotes the duality between V_n and V_n^* . Then it is easy to see that due to conditions **(1)**, **(2)** and **(3)** in Assumption 2.2, the operators A^n and B^n satisfy **(S1)**, **(S2)** and **(S3)** in Assumption 3.1, respectively. Furthermore, taking into account Remark 2.4 it is obvious that **(S4)** holds. Assume the regularity condition **(R3)** from Assumption 2.3. Then by virtue of the definition of Π_n , A^n and B^n , due to Lipschitz conditions **(3)** in Assumption 2.2 and (2.8) in Remark 2.4, we have, recalling the direct inequality (2.14),

$$\begin{aligned} & |\Pi_n A(t, u) - A^n(t, \Pi_n u)|_{V_n^*}^2 + \sum_k |\Pi_n B_k(t, u) - B_k^n(t, \Pi_n u)|_{H_n}^2 \\ & \leq |A(t, u) - A^n(t, \Pi_n u)|_{V^*}^2 + \sum_k |B_k(t, u) - B_k^n(t, \Pi_n u)|_H^2 \\ & \leq C(L_1 + L_2) 2^{-2nl} |u|_{\mathcal{V}}^2 \end{aligned}$$

almost surely for all $t \in [0, T]$ and $u \in \mathcal{V}$, which yields **(Cn)** with $\xi^n := 0$ and $\varepsilon_n := C(L_1 + L_2)2^{-nl}$. In the last section we will give examples of operators such that Assumption 2.3 holds.

4. IMPLICIT SPACE-TIME DISCRETIZATIONS

4.1. Description of the scheme. For a fixed integer $m \geq 1$ set $\tau := T/m$ and $t_i = i\tau$ for $i = 0, \dots, m$. Let $V_n \hookrightarrow H_n \hookrightarrow V_n^*$ satisfy the conditions in section 2.2. Given a V_n -valued \mathcal{F}_0 -measurable random variable $u_0^{n,\tau}$ and $\mathcal{F}_{t_i} \otimes \mathcal{B}(V_n)$ -measurable mappings

$$A_j^{n,\tau} : \Omega \times V_n \rightarrow V_n^* \quad \text{and} \quad B_{k,i}^{n,\tau} : \Omega \times V_n \rightarrow H_n, \quad \text{for } k = 1, \dots, d_1,$$

$j = 1, \dots, m$ and $i = 0, \dots, m-1$, consider for each n the system of equations

$$u_{i+1}^{n,\tau} = u_i^{n,\tau} + \tau A_{i+1}^{n,\tau}(u_{i+1}^{n,\tau}) + \sum_k B_{k,i}^{n,\tau}(u_i^{n,\tau})(W^k(t_{i+1}) - W^k(t_i)), \quad (4.1)$$

$i = 0, \dots, m-1$, for V_n -valued \mathcal{F}_{t_i} -measurable random variables $u_i^{n,\tau}$, $i = 1, \dots, m$.

Assumption 4.1. For almost all $\omega \in \Omega$ the operators $A_j^{n,\tau}$ and $B_{k,i}^{n,\tau}$ satisfy the following conditions for all $j = 1, \dots, m$, $i = 0, \dots, m-1$,

(ST1) (Strong monotonicity) There exist constants $\lambda > 0$ and $L \geq 0$ such that

$$\begin{aligned} 2\langle u - v, A_j^{n,\tau}(u) - A_j^{n,\tau}(v) \rangle_n + \sum_k |B_{k,j}^{n,\tau}(u) - B_{k,j}^{n,\tau}(v)|_{H_n}^2 \\ \leq -\lambda \|u - v\|_{V_n}^2 + L \|u - v\|_{H_n}^2 \end{aligned} \quad (4.2)$$

for all $u, v \in V_n$, $m \geq 1$, $n \geq 0$.

(ST2) (Growth condition on $A_i^{n,\tau}$ and $B_i^{n,\tau}$) There is a constant K such that

$$|A_j^{n,\tau}(u)|_{V_n^*}^2 \leq K \|u\|_{V_n}^2 + f_j^{n,\tau}, \quad \sum_k |B_{k,i}^{n,\tau}(u)|_{H_n}^2 \leq K \|u\|_{V_n}^2 + g_i^{n,\tau}$$

for all $u \in V_n$, $m \geq 1$, $n \geq 0$, where $f_j^{n,\tau}$ and $g_i^{n,\tau}$ are non-negative random variables, such that

$$\sup_{n,m} \sum_j \tau E f_i^{n,\tau} \leq M < +\infty, \quad \sup_{n,m} \max_i E g_i^{n,\tau} \leq M < +\infty.$$

(ST3) (Lipschitz condition on $A_j^{n,\tau}$) There exists a constants L_1 such that

$$|A_j^{n,\tau}(u) - A_j^{n,\tau}(v)|_{V_n^*}^2 \leq L_1 \|u - v\|_{V_n}^2 \quad (4.3)$$

for all $u, v \in V_n$, $m \geq 1$, $n \geq 0$.

Remark 4.1. Clearly, conditions (ST1) and (ST3) imply the Lipschitz continuity of $B_{k,i}^{n,\tau}$ in $v \in V_n$, i.e., there is a constant $L_2 = L_2(L, \lambda, L_1)$ such that almost surely

$$\sum_k |B_{k,j}^{n,\tau}(u) - B_{k,j}^{n,\tau}(v)|_{H_n}^2 \leq L_2 \|u - v\|_{V_n}^2 \quad (4.4)$$

for all $u, v \in V_n$, $n \geq 1$, $m \geq 0$ and $j = 1, \dots, m$.

Remark 4.2. Conditions (ST1)–(ST2) imply that almost surely

$$2\langle u, A_j^{n,\tau}(u) \rangle_n + \sum_k |B_{k,j}^{n,\tau}(u)|_{H_n}^2 \leq -\frac{\lambda}{2} \|u\|_{V_n}^2 + L \|u\|_{H_n}^2 + C(f_j^{n,\tau} + g_j^{n,\tau})$$

for all $u \in V_n$, $n \geq 0$, $m \geq 1$ and $j = 1, \dots, m$, where $C = C(\lambda, K)$ is a constant. The Lipschitz condition (ST3) obviously implies that $A_i^{n,\tau}$ is hemicontinuous.

Proposition 4.3. Let Assumption 4.1 hold. Assume $E\|u_0^{n,\tau}\|_{V_n}^2 < \infty$ for all $n \geq 0$ and $m \geq 1$. Then for $\tau < 1/L$ equation (4.1) has a unique V_n -valued solution $(u_i^{n,\tau})_{j=1}^m$, such that $u_j^{n,\tau}$ is \mathcal{F}_{t_j} -measurable and $E\|u_j^{n,\tau}\|_{V_n}^2$ is finite for each j , n . (Here $1/L := \infty$ if $L = 0$.)

Proof. Equation (4.1) can be rewritten as

$$D_{i+1}(u_{i+1}^{n,\tau}) = u_i^{n,\tau} + \sum_k B_{k,i}^{n,\tau}(u_i^{n,\tau}) (W^k(t_{i+1}) - W^k(t_i)), \quad (4.5)$$

where $D_i : V_n \rightarrow V_n^*$ is defined by $D_i(v) = v - \tau A_i^{n,\tau}(v)$ for each $i = 1, 2, \dots, m$. Due to Assumption 4.1 and Remark 4.2 the operator D_i satisfies the assumptions (monotonicity, coercivity, linear growth and hemicontinuity) of Proposition 3.4 in [7] with $p = 2$. By virtue of this proposition, for $\tau < 1/L$, equation (4.5) has a unique

V_n -valued $\mathcal{F}_{t_{i+1}}$ -measurable solution $u_{i+1}^{n,\tau}$ for every given V -valued \mathcal{F}_{t_i} -measurable random variable $u_i^{n,\tau}$, and

$$\begin{aligned} E\|u_{i+1}^{n,\tau}\|_{V_n}^2 &\leq C E\left(1 + f_i^{n,\tau} + g_i^{n,\tau} + \left|\sum_k B_{k,i}^{n,\tau}(u_i^{n,\tau})(W^k(t_{i+1}) - W^k(t_i))\right|^2\right) \\ &\leq C\left(1 + E f_i^{n,\tau} + E g_i^{n,\tau} + \sum_k \tau E|B_{k,i}^{n,\tau}(u_i^{n,\tau})|^2_n\right) \\ &\leq C(1 + E f_i^{n,\tau} + E g_i^{n,\tau} + K\tau E\|u_i^{n,\tau}\|_{V_n}^2 + \tau E g_i^{n,\tau}), \end{aligned}$$

where $C = C(\lambda, \tau)$ is a constant. Hence induction on i concludes the proof. \square

4.2. Rate of convergence of the implicit scheme. Let Assumption 2.3 on the regularity of equation 2.1 and its solution u hold. We relate the operators $A(t_i, \cdot)$ and $A_i^{n,\tau}$ as well as the operators $B_k(t_i, \cdot)$ and $B_{k,i}^{n,\tau}$ by the following consistency assumption.

Condition (Cn τ) (*Consistency*) *There exist constants $\nu \in]0, \frac{1}{2}]$, $c \geq 0$, a sequence of numbers $\varepsilon_n \rightarrow 0$, such that almost surely*

$$\begin{aligned} |\Pi_n A(t_j, u) - A_j^{n,\tau}(\Pi_n u)|_{V_n^*}^2 &\leq c(|u|_{\mathcal{V}}^2 + \xi_j^{n,\tau})(\tau^{2\nu} + \varepsilon_n^2), \\ \sum_k |\Pi_n B_k(t_i, u) - B_{k,i}^{n,\tau}(\Pi_n u)|_{H_n}^2 &\leq c(|u|_{\mathcal{V}}^2 + \eta_i^{n,\tau})(\tau^{2\nu} + \varepsilon_n^2) \end{aligned}$$

for all $j = 1, \dots, m$, $i = 0, \dots, m-1$ and $u \in \mathcal{V}$, where $\xi_j^{n,\tau}$ and $\eta_i^{n,\tau}$ are non-negative random variables such that

$$\sup_{n,m} \sum_j \tau E \xi_j^{n,\tau} \leq M, \quad \sup_{n,m} \sum_i \tau E \eta_i^{n,\tau} \leq M.$$

Theorem 4.4. *Let Assumptions 2.3 and 4.1 as well as condition (Cn τ) hold. Assume*

$$\sup_{n,m} E\|u_0^{n,\tau}\|_{V_n}^2 \leq M. \quad (4.6)$$

Set $e_i^{n,\tau} = \Pi_n u(t_i) - u_i^{n,\tau}$. Then for $\tau < 1/L$ and $n \geq 0$

$$\begin{aligned} E \max_{0 \leq i \leq m} |e_i^{n,\tau}|_{H_n}^2 + \sum_{1 \leq i \leq m} \tau E\|e_i^{n,\tau}\|_{V_n}^2 \\ \leq C_1 E|e_0^{n,\tau}|_{H_n}^2 + C_2(\tau^{2\nu} + \varepsilon_n^2)(r_1 + r_2 + M), \end{aligned} \quad (4.7)$$

where $C_1 = C_1(\lambda, L, T)$ and $C_2 = C_2(\lambda, L, K, T, p, c, L_1, L_2)$ are constants.

Proof. We fix n, τ , and to ease notation we write e_i, A_i and $B_{k,i}$ in place of $e_i^{n,\tau}, A_i^{n,\tau}$ and $B_{k,i}^{n,\tau}$, respectively. Similarly, we often use u_i in place of $u_i^{n,\tau}$ for $i = 1, 2, \dots, m$. Then for any $i = 0, \dots, m-1$,

$$\begin{aligned} |e_{i+1}|_n^2 - |e_i|_n^2 &= 2 \int_{t_i}^{t_{i+1}} \langle e_{i+1}, \Pi_n A(s, u(s)) - A_{i+1}(u_{i+1}) \rangle_n ds \\ &\quad + 2 \sum_k \int_{t_i}^{t_{i+1}} (e_{i+1}, F_k(s))_n dW^k(s) \\ &\quad - \left| \int_{t_i}^{t_{i+1}} [\Pi_n A(s, u(s)) - A_{i+1}(u_{i+1})] ds + \sum_k \int_{t_i}^{t_{i+1}} F_k(s) dW^k(s) \right|_n^2 \end{aligned}$$

$$\begin{aligned}
 &= 2 \int_{t_i}^{t_{i+1}} \langle e_{i+1}, \Pi_n A(s, u(s)) - A_{i+1}(u_{i+1}) \rangle_n ds \\
 &\quad + \left| \sum_k \int_{t_i}^{t_{i+1}} F_k(s) dW^k(s) \right|_n^2 + 2 \sum_k \int_{t_i}^{t_{i+1}} (e_i, F_k(s))_n dW^k(s) \\
 &\quad - \left| \int_{t_i}^{t_{i+1}} [\Pi_n A(s, u(s)) - A_{i+1}(u_{i+1})] ds \right|_n^2,
 \end{aligned}$$

where for $k = 1, \dots, d_1$ one sets

$$F_k(s) = \Pi_n B_k(s, u(s)) - B_{k,i}(u_i^{n,\tau}), \quad s \in]t_i, t_{i+1}], \quad i = 0, 1, \dots, m-1.$$

Summing up for $i = 0, \dots, l-1$, we obtain

$$|e_l|_n^2 \leq |e_0|_n^2 + 2 \sum_{0 \leq i < l} \int_{t_i}^{t_{i+1}} \langle e_{i+1}, \Pi_n A(s, u(s)) - A_{i+1}(u_{i+1}) \rangle_n ds + Q(t_l) + I(t_l), \quad (4.8)$$

where

$$\begin{aligned}
 Q(t_l) &= \sum_{0 \leq i < l} \left| \sum_k \int_{t_i}^{t_{i+1}} F_k(s) dW^k(s) \right|_n^2, \\
 I(t_l) &= 2 \sum_k \int_0^{t_l} (e(s), F_k(s))_n dW^k(s), \quad e(s) := e_i \text{ for } s \in]t_i, t_{i+1}], \quad i = 0, \dots, m.
 \end{aligned}$$

First we show

$$\sup_{0 \leq l \leq m} E|e_l|_n^2 + E \sum_{1 \leq i \leq m} \tau \|e_i\|_n^2 \leq C_1 E|e_0|_n^2 + C_2 (\tau^{2\nu} + \varepsilon_n^2) (r_1 + r_2 + M), \quad (4.9)$$

where $C_1 = C_1(\lambda, L, T)$ and $C_2 = C_2(\lambda, L, K, T, p, c, L_1, L_2)$ are constants. To this end we take expectation in both sides of (4.8) and use the strong monotonicity condition **(ST1)** from Assumption 4.1 to get

$$\begin{aligned}
 E|e_l|_n^2 &\leq E|e_0|_n^2 + 2E \sum_{0 \leq i < l} \tau \langle e_{i+1}, A_{i+1}(\Pi_n u(t_{i+1})) - A_{i+1}(u_{i+1}) \rangle_n \\
 &\quad + E \sum_{0 \leq i < l-1} \sum_k \tau |B_{k,i+1}(\Pi_n u(t_{i+1})) - B_{k,i+1}(u_{i+1})|_n^2 + \sum_{1 \leq j \leq 3} S_j \\
 &\leq E|e_0|_n^2 - \lambda \sum_{1 \leq i \leq l} \tau E \|e_i\|_n^2 + L \sum_{1 \leq i \leq l} \tau E |e_i|_n^2 + \sum_{1 \leq j \leq 3} S_j \quad (4.10)
 \end{aligned}$$

for $l = 1, \dots, m$, where

$$\begin{aligned}
 S_1 &= 2 \sum_{1 \leq i \leq l} E \int_{t_{i-1}}^{t_i} \langle e_i, \Pi_n A(s, u(s)) - A_i(\Pi_n u(t_i)) \rangle_n ds, \\
 S_2 &= \sum_k \sum_{1 \leq i < l} E \int_{t_i}^{t_{i+1}} [|F_k(s)|_n^2 - |B_{k,i}(\Pi_n u(t_i)) - B_{k,i}(u_i)|_n^2] ds, \\
 S_3 &= \sum_k E \int_0^\tau |F_k(s)|_n^2 ds.
 \end{aligned}$$

For any $\varepsilon > 0$

$$S_1 \leq \varepsilon \sum_{1 \leq i \leq l} \tau E \|e_i\|_n^2 + \frac{1}{\varepsilon} R,$$

where

$$\begin{aligned} R &= R(t_l) = \sum_{1 \leq i \leq l} E \int_{t_{i-1}}^{t_i} |\Pi_n A(s, u(s)) - A_i(\Pi_n u(t_i))|_{n^*}^2 ds \leq 3 \sum_{1 \leq j \leq 3} R_j, \quad (4.11) \\ R_1 &= \sum_{1 \leq i \leq l} E \int_{t_{i-1}}^{t_i} |\Pi_n A(s, u(s)) - \Pi_n A(t_i, u(s))|_{n^*}^2 ds, \\ R_2 &= \sum_{1 \leq i \leq l} E \int_{t_{i-1}}^{t_i} |\Pi_n A(t_i, u(s)) - A_i(\Pi_n u(s))|_{n^*}^2 ds, \\ R_3 &= \sum_{1 \leq i \leq l} E \int_{t_{i-1}}^{t_i} |A_i(\Pi_n u(s)) - A_i(\Pi_n u(t_i))|_{n^*}^2 ds. \end{aligned}$$

Due to condition (2.11) on the time regularity of A in Assumption 2.3, (2.13), **(Cn τ)**, the Lipschitz condition (4.3) in Assumption 4.1 and inequality (2.12) from Remark 2.5, we deduce

$$R_1 \leq \tau^{2\nu} p^2 E \int_0^T (K |u(s)|_{\mathcal{V}}^2 + \eta) ds, \quad (4.12)$$

$$R_2 \leq c(\tau^{2\nu} + \varepsilon_n^2) \left(E \int_0^T |u(s)|_{\mathcal{V}}^2 ds + \sum_{1 \leq i \leq m} \tau E \xi_i^{n,\tau} \right), \quad (4.13)$$

$$R_3 \leq L_1 p^2 \sum_{1 \leq i \leq l} \int_{t_{i-1}}^{t_i} E \|u(s) - u(t_i)\|^2 ds \leq T L_1 p^2 M_1 \tau, \quad (4.14)$$

with $M_1 := C(r_1 + r_2 + M)$. By (2.13), the regularity condition **(R3)** on B from Assumption 2.3, the growth condition **(ST2)** on $B_{i,k}$ from Assumption 4.1, and by condition (4.6) on the initial values we have

$$\begin{aligned} S_3 &\leq 2 \sum_k \int_0^\tau E |\Pi_n B_k(s, u(s))|_n^2 ds + 2 \sum_k \tau E |B_{k,0}(u_0^{n,\tau})|_n^2 \\ &\leq 2\tau p^2 \left(K \sup_{t \in [0, T]} E \|u(t)\|_{\mathcal{H}}^2 + \sup_{t \in [0, T]} E \eta(t) \right) \\ &\quad + 2\tau \left(K \sup_{n,m} E \|u_0^{n,\tau}\|_n^2 + \sup_{n,m} E g_0^{n,\tau} \right). \quad (4.15) \end{aligned}$$

Using the simple inequality $|b|_n^2 - |a|_n^2 \leq \varepsilon |a|_n^2 + (1 + \frac{1}{\varepsilon}) |b - a|_n^2$ with

$$a := B_{k,i}(\Pi_n u(t_i)) - B_{k,i}(u_i), \quad b := F_k(s),$$

for any $\varepsilon > 0$ we have $S_2 \leq \varepsilon P_1 + (1 + \frac{1}{\varepsilon}) P_2$ with

$$\begin{aligned} P_1 = P_{1l} &= \sum_{1 \leq i \leq l} E \sum_k \tau |B_{k,i}(\Pi_n u(t_i)) - B_{k,i}(u_i)|_n^2, \\ P_2 = P_{2l} &= \sum_{1 \leq i < l} E \int_{t_i}^{t_{i+1}} \sum_k |\Pi_n B_k(s, u(s)) - B_{k,i}(\Pi_n u(t_i))|_n^2 ds. \quad (4.16) \end{aligned}$$

By Remark 4.1 on the Lipschitz continuity of $B_{k,i}$ we get $P_1 \leq L_2 E \sum_{1 \leq i \leq l} \tau \|e_i\|_n^2$. Clearly, $P_2 \leq 3(Q_1 + Q_2 + Q_3)$ with

$$\begin{aligned} Q_1 &= \sum_{1 \leq i < l} E \int_{t_i}^{t_{i+1}} \sum_k |\Pi_n B_k(s, u(s)) - \Pi_n B_k(t_i, u(s))|_n^2 ds, \\ Q_2 &= \sum_{1 \leq i < l} E \int_{t_i}^{t_{i+1}} \sum_k |\Pi_n B_k(t_i, u(s)) - B_{k,i}(\Pi_n u(s))|_n^2 ds, \\ Q_3 &= \sum_{1 \leq i < l} E \int_{t_i}^{t_{i+1}} \sum_k |B_{k,i}(\Pi_n u(s)) - B_{k,i}(\Pi_n u(t_i))|_n^2 ds. \end{aligned}$$

Due to **(R4)** (ii) in Assumption 2.3 on the time regularity of B , consistency **(Cn τ)**, the Lipschitz continuity of $B_{k,i}$ proved in Remark 4.1, (2.12) proved in Remark 2.5 and (2.13),

$$\begin{aligned} Q_1 &\leq \tau^{2\nu} p^2 \left(K E \int_0^T |u(s)|_V^2 ds + T E \eta \right), \\ Q_2 &\leq c(\tau^{2\nu} + \varepsilon_n^2) \left(E \int_0^T |u(s)|_V^2 ds + \sup_{n,m} \sum_{0 \leq i < l} \tau E \eta_i^{n,\tau} \right), \\ Q_3 &\leq L_2 p^2 T \sup_{|t-s| \leq \tau} E \|u(t) - u(s)\|^2 \leq \tau L_2 p^2 T M_1. \end{aligned}$$

Hence

$$\begin{aligned} S_2 &\leq \varepsilon L_2 E \sum_{1 \leq i \leq l} \tau \|e_i\|_n^2 + C \left(1 + \frac{1}{\varepsilon} \right) (\tau^{2\nu} + \varepsilon_n^2) \\ &\quad \times \left(E \int_0^T |u(s)|_V^2 ds + T E \eta + \sup_{n,m} \sum_i \tau E \eta_i^{n,\tau} + M_1 \right), \end{aligned} \quad (4.17)$$

where $C = C(p, K, L_2, c)$. Choosing $\varepsilon > 0$ sufficiently small, from (4.10) and (4.12)–(4.17) we obtain for $l = 1, \dots, m$,

$$\begin{aligned} E|e_l|_n^2 + \frac{\lambda}{2} E \sum_{1 \leq i \leq l} \tau \|e_i\|_n^2 \\ \leq E|e_0|_n^2 + L \sum_{1 \leq i \leq l} \tau E|e_i|_n^2 + C(\tau^{2\nu} + \varepsilon_n^2)(r_1 + r_2 + M), \end{aligned} \quad (4.18)$$

where $C = C(K, \lambda, p, T, c, L_1, L_2)$ is a constant. Since $\sup_m \sum_{i=1}^m \tau = T < +\infty$, if $L\tau < 1$ a discrete version of Gronwall's lemma yields the existence of constants $C_1 = C_1(L, \lambda, T)$ and $C_2 = C_2(L, K, \lambda, p, T, c, L_1, L_2)$ such that for sufficiently large m

$$\max_{1 \leq l \leq m} E|e_l|_n^2 \leq C_1 E|e_0|_n^2 + C_2(r_1 + r_2 + M)(\tau^{2\nu} + \varepsilon_n^2)$$

holds for all n . This together with (4.18) concludes the proof of (4.9). To prove (4.7) notice that from (4.8) by the same calculations as above, but taking first max

in l and then expectation, we get

$$\begin{aligned} E \max_{1 \leq i \leq m} |e_i|_n^2 &\leq E|e_0|_n^2 + E \sum_{1 \leq i \leq m} \tau \|e_i\|_n^2 + L \sum_{1 \leq i \leq l} \tau E|e_i|_n^2 \\ &\quad + R(T) + R_0 + E \max_{0 \leq i \leq m} I(t_i) + EQ(T). \end{aligned} \quad (4.19)$$

where $R(T) = R(t_m)$ is defined by (4.11) and

$$R_0 = P_{1m} + 2P_{2m} + S_3. \quad (4.20)$$

The terms $R(T)$, P_{1m} and P_{2m} have already been estimated above by the right-hand side of (4.18) and S_3 has been estimated by (4.15). Notice that

$$EQ(T) = E \int_0^T \sum_k |F_k(s)|_n^2 ds \leq 2P_{1m} + 2P_{2m},$$

and by Davis' inequality

$$\begin{aligned} E \max_{0 \leq i \leq m} I(t_i) &\leq 6E \left\{ \int_0^T \sum_k |(e(s), F_k(s))_n|^2 ds \right\}^{1/2} \\ &\leq \frac{1}{2} E \max_{0 \leq i \leq m} |e_i|_n^2 + 18E \int_0^T \sum_k |F_k(s)|_n^2 ds. \end{aligned}$$

Thus from (4.19) we obtain (4.7). \square

Remark 4.5. *One can show, like it is observed in [8], that if instead of the Lipschitz condition (4.3) we assume that $A_i^{n,\tau}$ are hemicontinuous and $B_{k,i}^{n,\tau}$ satisfy the Lipschitz condition (4.4), then the order of the speed of convergence is divided by two.*

4.3. Examples. (i) Consider from Example 3.3 the normal triples

$$V \hookrightarrow H^* \hookrightarrow V^*, \quad V_n \hookrightarrow H_n^* \hookrightarrow V_n^*$$

with the orthogonal projection $\Pi_n : H = L^2(\mathbb{R}^d) \rightarrow H_n$ and auxiliary spaces $\mathcal{H} = W_2^{r+\rho}(\mathbb{R}^d)$ and $\mathcal{V} = W_2^{r+l}(\mathbb{R}^d)$ for some $l > \rho \geq 0$.

Let A and $B = (B_k)$ be $\mathcal{P} \otimes \mathcal{B}(V)$ -measurable mappings from $[0, \infty[\times \Omega \times V$ into V^* and H^{d_1} , respectively, satisfying Assumptions 2.2 and 2.3 such that f and g in (2.6) satisfies

$$\sup_{t \in [0, T]} f(t) \leq M, \quad \sup_{t \in [0, T]} g(t) \leq M.$$

For $\omega \in \Omega$, $j = 1, \dots, m$ and $i = 0, \dots, m-1$ let $A_j^{n,\tau}(\omega, \cdot) : V^n \rightarrow V_n^*$ and $B_{k,i}^{n,\tau}(\omega, \cdot) : V^n \rightarrow H_n$ be defined by

$$\langle A_j^{n,\tau}(\omega, u), v \rangle_n = \langle A(t_j, \omega, u), v \rangle \quad \text{and} \quad B_{k,i}^{n,\tau}(\omega, u) = \Pi_n B_k(t_i, \omega, u) \quad (4.21)$$

for all $u, v \in V_n$, where $\langle \cdot, \cdot \rangle_n$ denotes the duality between V_n and V_n^* . Then it is easy to see, like in Example 3.3, that due to **(1)**, **(2)** and **(3)** in Assumption 2.2, **(ST1)**, **(ST2)** and **(ST3)** in Assumption 4.1 hold respectively. In the same way as **(Cn)** is verified in Example 3.3, one can also easily show that the consistency assumption **(Cn τ)** holds.

(ii) Another choice for $A_j^{n\tau}$ and $B_{k,i}^{n\tau}$ can be defined by

$$\begin{aligned} \langle A_j^{n,\tau}(u), v \rangle_n &= \frac{1}{\tau} \int_{t_{j-1}}^{t_j} \langle A(s, u), v \rangle ds, \quad u, v \in V_n, \\ B_{k,0}^{n,\tau}(u) &= \Pi_n B_k(0, u), \quad B_{k,j}^{n,\tau}(u) = \frac{1}{\tau} \int_{t_{j-1}}^{t_j} \Pi_n B_k(s, u) ds, \quad u \in V_n, \end{aligned} \quad (4.22)$$

instead of (4.21). One can show by a similar computation as before, combined with the use of Jensen's inequality, that Assumption 2.2 and **(R3)**-**(R4)** in Assumption 2.3 imply Assumption 4.1 and condition **(Cn τ)**.

(iii) Finally, let $V_n = V$, $H_n = H$ and let Π_n be the identity operator for every n . Let Assumptions 2.2 and 2.3 hold. Then one recovers the conclusions of Theorems 3.2 and 3.4 in [8] concerning the rate of convergence of the implicit time discretization scheme with $\varepsilon_n = 0$.

5. EXPLICIT SPACE-TIME DISCRETIZATION SCHEME

5.1. Description of the scheme. Let V_n , H_n and V_n^* be a normal triple and Π_n be continuous linear operators which satisfy the condition (2.13). Assume moreover that for each $n \geq 0$ as sets

$$V_n = H_n = V_n^*,$$

and there is a constant $\vartheta(n)$ such that

$$\|u\|_{V_n}^2 \leq \vartheta(n) |u|_{H_n}^2, \quad \forall u \in H_n. \quad (5.1)$$

Then by duality we also have

$$|u|_{H_n}^2 \leq \vartheta(n) |u|_{V_n^*}^2, \quad \forall u \in V_n^*.$$

Consider for each n and $i = 0, 1, \dots, m-1$ the equations

$$u_{\tau,i+1}^n = u_{\tau,i}^n + \tau A_i^{n,\tau}(u_{\tau,i}^n) + \sum_k B_{k,i}^{n,\tau}(u_{\tau,i}^n) (W^k(t_{i+1}) - W^k(t_i)), \quad (5.2)$$

for V_n -valued \mathcal{F}_{t_i} -measurable random variables $u_{\tau,i}^n$ for $i = 1, \dots, m$, where $u_{\tau,0}^n$ is a given V_n -valued \mathcal{F}_0 -measurable random variable, and

$$A_i^{n,\tau} : \Omega \times V_n \rightarrow V_n^* \quad \text{and} \quad B_{k,i}^{n,\tau} : \Omega \times V_n \rightarrow H_n$$

are given $\mathcal{F}_{t_i} \otimes \mathcal{B}(V_n)$ -measurable mappings such that Assumption 4.1 holds.

Proposition 5.1. *Let Assumption 4.1 hold. Then for any V -valued \mathcal{F}_0 -measurable random variable $u_{\tau,0}^n$ such that $E\|u_{\tau,0}^n\|_{V_n}^2 < \infty$, the system of equations (5.2) has a unique solution $(u_{\tau,i}^n)_{i=1}^m$ such that $u_{\tau,i}^n$ is \mathcal{F}_{t_i} -measurable and $E\|u_{\tau,i}^n\|_{V_n}^2 < \infty$ for all i , m and n .*

Proof. By (5.1) we have $\|u_{\tau,i+1}^n\|_n^2 \leq \vartheta(n) |u_{\tau,i+1}^n|_n^2$, and by (5.2)

$$\begin{aligned} E|u_{\tau,i+1}^n|_n^2 &\leq 3E|u_{\tau,i}^n|_n^2 + 3\tau E|A_i^{n,\tau}(u_{\tau,i}^n)|_n^2 + 3\tau \sum_k E|B_{k,i}^{n,\tau}(u_{\tau,i}^n)|_n^2 \\ &\leq 3(\vartheta(n) + \vartheta(n)\tau K + \tau K) E\|u_{\tau,i}^n\|_n^2 + 3\tau(\vartheta(n) + 1)M. \end{aligned}$$

Hence we get the proposition by induction on i . \square

5.2. Rate of convergence of the scheme. The following theorem gives the rate of convergence of $e_{\tau,i}^n := \Pi_n u(t_i) - u_{\tau,i}^n$.

Theorem 5.2. *Let Assumption 2.3, Assumption 4.1 with index $j = i = 0, \dots, m-1$ in its formulation, and the consistency condition $(\mathbf{Cn}\tau)$ hold. Let n and τ satisfy*

$$L_1 \tau \vartheta(n) + 2\sqrt{L_1 L_2 \tau \vartheta(n)} \leq q \quad (5.3)$$

for some constant $q < \lambda$, where L_1 and L_2 are the Lipschitz constants in (4.3) and (4.4), respectively. Then

$$E \max_{0 \leq i \leq m} |e_{\tau,i}^n|_{H_n}^2 + \sum_{0 \leq i < m} \tau E \|e_{\tau,i}^n\|_{V_n}^2 \leq C_1 E |e_{\tau,0}^n|_{H_n}^2 + C_2 (\tau^{2\nu} + \varepsilon_n^2) (r_1 + r_2 + M), \quad (5.4)$$

where $C_1 = C_1(\lambda, q, L, T)$ and $C_2 = C_2(\lambda, q, L, K, T, p, c, L_1, L_2)$ are constants.

Proof. Note that when we refer to any condition in Assumption 4.1 then we mean it with the index j replaced in its formulation with i running through $0, \dots, m-1$. To ease notation we omit the indices n and τ from $e_{\tau,i}^n$, $u_{\tau,i}^n$, $A_i^{n,\tau}$ and $B_i^{n,\tau}$ when this does not cause ambiguity. For any $i = 0, \dots, m-1$

$$\begin{aligned} |e_{i+1}|_n^2 - |e_i|_n^2 &= 2 \int_{t_i}^{t_{i+1}} \langle e_i, \Pi_n A(s, u(s)) - A_i(u_i) \rangle_n ds \\ &+ 2 \sum_k \int_{t_i}^{t_{i+1}} (e_i, F_k(s))_n dW^k(s) + \sum_k \left| \int_{t_i}^{t_{i+1}} F_k(s) dW^k(s) \right|_n^2 \\ &+ \left| \int_{t_i}^{t_{i+1}} [\Pi_n A(s, u(s)) - A_i(u_i)] ds \right|_n^2 \\ &+ 2 \sum_k \left(\int_{t_i}^{t_{i+1}} [\Pi_n A(s, u(s)) - A_i(u_i)] ds, \int_{t_i}^{t_{i+1}} F_k(s) dW^k(s) \right)_n, \end{aligned}$$

where

$$F_k(s) = \Pi_n B_k(s, u(s)) - B_{k,i}(u_i), \quad s \in]t_i, t_{i+1}], \quad i = 0, 1, \dots, m-1.$$

Hence for $l = 1, \dots, m$ and every $\delta > 0$,

$$\begin{aligned} |e_l|_n^2 &\leq |e_0|_n^2 + 2 \sum_{0 \leq i < l} \int_{t_i}^{t_{i+1}} \langle e_i, \Pi_n A(s, u(s)) - A_i(u_i) \rangle_n ds + 2I(t_l) + Q(t_l) \\ &+ (1 + \frac{1}{\delta})S(t_l) + \delta Q(t_l), \end{aligned} \quad (5.5)$$

where

$$\begin{aligned} I(t_l) &= \int_0^{t_l} (e(s), F_k(s)) dW^k(s), \quad e(s) := e_i \quad \text{for } s \in]t_i, t_{i+1}], \quad i \geq 0, \\ S(t_l) &= \sum_{0 \leq i < l} \left| \int_{t_i}^{t_{i+1}} [\Pi_n A(s, u(s)) - A_i(u_i)] ds \right|_n^2, \\ Q(t_l) &= \sum_{0 \leq i < l} \sum_k \left| \int_{t_i}^{t_{i+1}} F_k(s) dW^k(s) \right|_n^2. \end{aligned}$$

First we prove

$$\max_{1 \leq i \leq m} E|e_i|_n^2 + \sum_{0 \leq i < m} \tau E \|e_i\|_n^2 \leq C_1 E|e_0|_n^2 + C_2(\tau^{2\nu} + \varepsilon_n^2)(r_1 + r_2 + M), \quad (5.6)$$

with some constants $C_1 = C_1(\lambda, q, L, T)$ and $C_2 = C_2(\lambda, q, L, K, T, p, c, L_1, L_2)$. To this end we take expectation in both sides of (5.5) and use the strong monotonicity condition (4.2) in Assumption 4.1, to get

$$\begin{aligned} E|e_l|_n^2 &\leq E|e_0|_n^2 + 2E \sum_{0 \leq i < l} \tau \langle e_i, A_i(\Pi_n u(t_i)) - A_i(u_i) \rangle_n \\ &\quad + E \sum_{0 \leq i < l} \tau |B_{k,i}(\Pi_n u(t_i)) - B_{k,i}(u_i)|_n^2 + \sum_{i=1,2} S_i + (1 + \frac{1}{\delta})ES(t_l) + \delta EQ(t_l) \\ &\leq E|e_0|_n^2 - \lambda E \sum_{0 \leq i < l} \tau \|e_i\|_n^2 + LE \sum_{0 \leq i < l} \tau |e_i|_n^2 + \sum_{i=1,2} S_i + (1 + \frac{1}{\delta})ES(t_l) + \delta EQ(t_l), \end{aligned}$$

for any $\delta > 0$, where

$$\begin{aligned} S_1 &= 2 \sum_{0 \leq i < l} E \int_{t_i}^{t_{i+1}} \langle e_i, \Pi_n A(s, u(s)) - A_i(\Pi_n u(t_i)) \rangle_n ds, \\ S_2 &= \sum_k \sum_{0 \leq i < l} E \int_{t_i}^{t_{i+1}} [|F_k(s)|_n^2 - |B_{k,i}(\Pi_n u(t_i)) - B_{k,i}(u_i)|_n^2] ds. \end{aligned}$$

As in the proof of Theorem 4.4 we get for any $\varepsilon > 0$,

$$\begin{aligned} S_1 &\leq \varepsilon \sum_{0 \leq i < l} \tau E \|e_i\|_n^2 + \frac{1}{\varepsilon} C(r_1 + r_2 + M)(\tau^{2\nu} + \varepsilon_n^2), \\ S_2 &\leq L_2 \varepsilon \sum_{0 \leq i < l} \tau E \|e_i\|_n^2 + \frac{1}{\varepsilon} C(r_1 + r_2 + M)(\tau^{2\nu} + \varepsilon_n^2) \end{aligned}$$

with a constant $C = C(K, p, T, L_1, L_2, c)$. Notice that for any $\varepsilon > 0$,

$$ES(t_l) \leq \tau \vartheta(n) J(t_l),$$

$$J(t_l) := \sum_{0 \leq i < l} E \int_{t_i}^{t_{i+1}} |\Pi_n A(s, u(s)) - A_i(u_i)|_{n^*}^2 ds \leq (1 + \varepsilon)R_0 + (1 + \frac{1}{\varepsilon})R, \quad (5.7)$$

$$EQ(t_l) \leq (1 + \varepsilon)P_1 + (1 + \frac{1}{\varepsilon})P_2, \quad (5.8)$$

where

$$\begin{aligned} P_1 &:= \sum_{0 \leq i < l} E \sum_k |B_{k,i}(\Pi_n u(t_i)) - B_{k,i}(u_i)|_n^2 \tau \leq L_2 E \sum_{0 \leq i < l} \tau \|e_i\|_n^2 \\ P_2 &:= \sum_{0 \leq i < l} E \int_{t_i}^{t_{i+1}} \sum_k |\Pi_n B_k(s, u(s)) - B_{k,i}(\Pi_n u(t_i))|_n^2 ds, \\ R_0 &:= E \sum_{0 \leq i < l} \tau |A_i(\Pi_n u(t_i)) - A_i(u_i)|_{n^*}^2 \leq L_1 E \sum_{0 \leq i < l} \tau \|e_i\|_n^2, \\ R &:= E \sum_{0 \leq i < l} \int_{t_i}^{t_{i+1}} |\Pi_n A(s, u(s)) - A_i(\Pi_n u(t_i))|_{n^*}^2 ds, \end{aligned}$$

for any $l = 1, 2, \dots, m$. In the same way as in the proof of Theorem 4.4 we obtain

$$R \leq C(\tau^{2\nu} + \varepsilon_n^2)(r_1 + r_2 + M), \quad (5.9)$$

and that

$$P_2 \leq C'(\tau^{2\nu} + \varepsilon_n^2)(r_1 + r_2 + M), \quad (5.10)$$

where $C = C(K, p, c, L_1, T)$ and $C' = C'(K, p, c, L_2, T)$ are constants. Consequently,

$$\begin{aligned} E|e_l|_n^2 &\leq E|e_0|_n^2 + (\mu - \lambda)E \sum_{0 \leq i < l} \tau \|e_i\|_n^2 + LE \sum_{0 \leq i < l} \tau |e_i|_n^2 \\ &\quad + (1 + \tau\vartheta(n))\left(1 + \frac{1}{\delta} + \frac{1}{\varepsilon}\right)C(\tau^{2\nu} + \varepsilon_n^2)(r_1 + r_2 + M) \end{aligned} \quad (5.11)$$

for any $\delta > 0$ and $\varepsilon > 0$, where

$$\mu = (1 + \varepsilon)\left[\left(1 + \frac{1}{\delta}\right)\tau\vartheta(n)L_1 + \delta L_2\right] + \varepsilon(1 + L_2),$$

and $C = C(K, p, c, T, L_1, L_2)$ is a constant. It is easy to see that due to (5.3)

$$\inf_{\delta > 0} \left(1 + \frac{1}{\delta}\right)\tau\vartheta(n)L_1 + \delta L_2 = \tau\vartheta(n)L_1 + 2\sqrt{\tau\vartheta(n)L_1 L_2} \leq q.$$

Therefore we can take $\delta > 0$ and $\varepsilon > 0$ such that $\mu \leq (q + \lambda)/2$. Thus from (5.11) we can get

$$\begin{aligned} E|e_l|_n^2 &\leq E|e_0|_n^2 - \frac{1}{2}(\lambda - q)E \sum_{0 \leq i < l} \tau \|e_i\|_n^2 + LE \sum_{0 \leq i < l} \tau |e_i|_n^2 \\ &\quad + C(\tau^{2\nu} + \varepsilon_n^2)(r_1 + r_2 + M), \end{aligned}$$

with a constant $C = C(K, \lambda, q, p, c, T, L_1, L_2)$. Hence by a discrete version of Gronwall's lemma we obtain (5.6). To prove (5.4) note that (5.5) yields

$$E \max_{1 \leq l \leq m} |e_l|_n^2 \leq |e_0|_n^2 + E \sum_{0 \leq i < l} \tau \|e_i\|_n^2 + 2ES(T) + 2E \max_{1 \leq l \leq m} I(t_l) + 2EQ(T), \quad (5.12)$$

where by (5.7)–(5.9) $ES(T) \leq \tau\vartheta(n)J(T)$, and

$$J(T) \leq 2L_1E \sum_{0 \leq i < m} \tau \|e_i\|_n^2 + 2C(\tau^{2\nu} + \varepsilon_n^2)(r_1 + r_2 + M).$$

By (5.8), (5.9) and (5.10)

$$EQ(T) \leq 2L_2 \sum_{0 \leq i < m} \tau \|e_i\|_n^2 + 2C'(\tau^{2\nu} + \varepsilon_n^2)(r_1 + r_2 + M).$$

Finally, in the same way as equation (4.21) is obtained, we get

$$\begin{aligned} E \max_{1 \leq i \leq m} I(t_i) &\leq 6E \left\{ \int_0^T \sum_k |(e(s), F_k(s))_n|^2 ds \right\}^{1/2} \\ &\leq \frac{1}{2}E \max_{0 \leq i < m} |e_i|_n^2 + 18E \int_0^T F_k^2(s) ds \leq \frac{1}{2}E \max_{0 \leq i < m} |e_i|_n^2 + 18EQ(T). \end{aligned}$$

Consequently, from (5.12) we obtain (5.4) by (5.6). \square

5.3. **Example.** Consider again the spaces

$$\mathcal{V} \subset \mathcal{H} \subset V \hookrightarrow H \hookrightarrow V^*, \quad V_n \hookrightarrow H_n \hookrightarrow V_n^*$$

from Examples 3.3 and 4.3. Notice that V_n , H_n and V_n^* are identified as sets and that due to the converse inequality (2.15) we have (5.1) with $\vartheta(n) = C2^{nr}$.

Let A and B satisfy the same conditions as in Example 4.3. Define $A_j^{n,\tau}$ and $B_{k,i}^{n,\tau}$ for $j = i = 0, \dots, m-1$ by (4.21) or define $A_j^{n,\tau}$ by (4.21) for $j = 0$ and $A_j^{n,\tau}$, $B_{k,i}^{n,\tau}$ by (4.22) for $j = 1, \dots, m-1$ and $i = 0, \dots, m-1$. Then, as shown in section 4.3, $A_j^{n,\tau}$ and $B_{k,i}^{n,\tau}$ satisfy the conditions in Assumption 4.1 as well as **(Cn τ)**. Hence, if the solution u satisfies **(R1)**-**(R2)** in Assumption 2.3, and $L_1\tau\vartheta_n + 2\sqrt{L_1L_2\tau\vartheta_n} \leq q < \lambda$, then the conditions of Theorem 5.2 hold.

6. EXAMPLES OF APPROXIMATIONS OF STOCHASTIC PDES

In this section we present some examples of stochastic PDEs for which the previous theorems provide rates of convergence for the above space and space-time discretization schemes. We refer to section 5 in [8] for more details. In this section for integers l the notation $|u|_l = |u|_{W^l}$ means the norm of u in $H^l = W^l(\mathbb{R}^d)$.

6.1. **Quasilinear equations.** Let us consider the stochastic partial differential equation

$$\begin{aligned} du(t, x) = & (Lu(t) + F(t, x, \nabla u(t, x), u(t, x))) dt \\ & + \sum_k (M_k u(t, x) + g_k(t, x)) dW^k(t), \quad t \in (0, T], x \in \mathbb{R}^d, \end{aligned} \quad (6.1)$$

with initial condition

$$u(0, x) = u_0(x), \quad x \in \mathbb{R}^d, \quad (6.2)$$

where F and g_k are Borel functions of $(\omega, t, x, p, r) \in \Omega \times [0, \infty) \times \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}$ and of $(\omega, t, x) \in \Omega \times [0, \infty) \times \mathbb{R}^d$, respectively, and L , M_k are differential operators of the form

$$L(t)v(x) = \sum_{|\alpha| \leq 1, |\beta| \leq 1} D^\alpha (a^{\alpha\beta}(t, x) D^\beta v(x)), \quad M_k(t)v(x) = \sum_{|\alpha| \leq 1} b_k^\alpha(t, x) D^\alpha v(x),$$

with functions $a^{\alpha\beta}$ and b_k^α of $(\omega, t, x) \in \Omega \times [0, \infty) \times \mathbb{R}^d$, for all multi-indices $\alpha = (\alpha_1, \dots, \alpha_d)$, $\beta = (\beta_1, \dots, \beta_d)$ of length $|\alpha| = \sum_i \alpha_i \leq 1$, $|\beta| \leq 1$. Here, and later on D^α denotes $D_1^{\alpha_1} \dots D_d^{\alpha_d}$ for any multi-indices $\alpha = (\alpha_1, \dots, \alpha_d) \in \{0, 1, 2, \dots\}^d$, where $D_i = \frac{\partial}{\partial x_i}$ and D_i^0 is the identity operator. We use the notation $\nabla_p := (\partial/\partial p_1, \dots, \partial/\partial p_d)$.

Let K and M denote some non-negative numbers. Fix an integer $l \geq 0$ and suppose that the following conditions hold:

Assumption (A1) (Stochastic parabolicity). *There exists a constant $\lambda > 0$ such that*

$$\sum_{|\alpha|=1, |\beta|=1} \left(a^{\alpha\beta}(t, x) - \frac{1}{2} \sum_k (b_k^\alpha b_k^\beta)(t, x) \right) z^\alpha z^\beta \geq \lambda \sum_{|\alpha|=1} |z^\alpha|^2$$

for all $\omega \in \Omega$, $t \in [0, T]$, $x \in \mathbb{R}^d$ and $z = (z^1, \dots, z^d) \in \mathbb{R}^d$, where $z^\alpha := z_1^{\alpha_1} z_2^{\alpha_2} \dots z_d^{\alpha_d}$ for $z \in \mathbb{R}^d$ and multi-indices $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_d)$.

Assumption (A2) (Smoothness of the initial condition). *Let u_0 be W_l^2 -valued \mathcal{F}_0 -measurable random variable such that $E|u_0|_l^2 \leq M$.*

Assumption (A3) (Smoothness of the linear term). *The derivatives of $a^{\alpha\beta}$ and b_k^α up to order l are $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable real functions such that almost surely*

$$|D^\gamma a^{\alpha\beta}(t, x)| + |D^\gamma b_k^\alpha(t, x)| \leq K, \quad \text{for all } |\alpha| \leq 1, |\beta| \leq 1, k = 1, \dots, d_1,$$

$t \in [0, T]$, $x \in \mathbb{R}^d$ and multi-indices γ with $|\gamma| \leq 2$.

Assumption (A4) (Smoothness of the nonlinear term). *The function F and their first order partial derivatives in p , x and r are $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d) \otimes \mathcal{B}(\mathbb{R}^d) \otimes \mathcal{B}(\mathbb{R})$ -measurable functions. The function g_k and its derivatives in x are $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable functions for every $k = 1, \dots, d_1$. There exists a constant K and a $\mathcal{P} \otimes \mathcal{B}$ -measurable function ξ of (ω, t, x) such that almost surely*

$$\begin{aligned} |\nabla_p F(t, x, p, r)| + \left| \frac{\partial}{\partial r} F(t, x, p, r) \right| &\leq K, \\ |F(t, \cdot, 0, 0)|_0^2 + \sum_k |g_k(t, \cdot)|_2^2 &\leq \eta, \\ |\nabla_x F(t, x, p, r)| &\leq L(|p| + |r|) + \xi(t, x), \quad |\xi(t)|_0^2 \leq \eta \end{aligned}$$

for all t, x, p, r , where η is a random variable such that $E\eta \leq M$.

Set $H = L^2(\mathbb{R}^d) = W_2^0$, $V = W_2^1$, $\mathcal{H} = W_2^2$ and $\mathcal{V} = W_2^3$ and suppose that the assumptions **(A1)**–**(A4)** hold with $l = 2$. Then the operators

$$A(t, \varphi) = L(t)\varphi + F(t, \cdot, \nabla\varphi, \varphi), \quad B_k(t, \varphi) = M_k(t)\varphi + g_k(t, \cdot), \quad \varphi \in V$$

and u_0 satisfy the conditions of Theorem 2.2. Hence (6.1)–(6.2) has a unique solution u on $[0, T]$. Furthermore, u has a W_2^2 -valued continuous modification such that

$$E \sup_{0 \leq t \leq T} |u(t)|_2^2 + E \int_0^T |u(t)|_3^2 dt < \infty.$$

Consequently the regularity conditions **(R1)** and **(R2)** in Assumption 2.3 hold. It is easy to check that A and B_k verify condition **(R3)**.

Assumption (A5) (Time regularity of A and B) *Almost surely*

(i)

$$\begin{aligned} \sum_k |D^\gamma (b_k^\alpha(t, x) - b_k^\alpha(s, x))|^2 &\leq K|t - s|, \\ \sum_k |g_k(s, \cdot) - g_k(t, \cdot)|_1^2 &\leq \eta|t - s|. \end{aligned}$$

(ii)

$$\begin{aligned} |D^\gamma (a^{\alpha,\beta}(t, x) - a^{\alpha,\beta}(s, x))|^2 &\leq K|t - s|, \\ |F(t, x, p, r) - F(s, x, p, r)|^2 &\leq K|t - s|(|p|^2 + |r|^2), \\ |\nabla_x F(t, x, p, r) - \nabla_x F(s, x, p, r)|^2 &\leq K|t - s|(|p|^2 + |r|^2), \\ |\nabla_p F(t, x, p, r) - \nabla_p F(s, x, p, r)|^2 &\leq K|t - s|, \\ \left| \frac{\partial}{\partial r} F(t, x, p, r) - \frac{\partial}{\partial r} F(s, x, p, r) \right|^2 &\leq K|t - s|. \end{aligned}$$

for all $|\alpha| \leq 1$, $|\beta| \leq 1$, $|\gamma| \leq 1$, $s, t \in [0, T]$ and $x \in \mathbb{R}^d$, where K is a constant and η is a random variable such that $E\eta \leq M$.

Clearly, Assumptions (i) and (ii) of **(A5)** imply conditions (i) and (ii) of **(R4)** in Assumption 2.3, respectively with $\nu = 1/2$.

Let H_n , V_n and Π_n be defined as in Example 2.6 and let $A^n(t, u)$ and $B_k^n(t, u)$ be defined by (3.12). Let $u_0 \in W_2^2 = \mathcal{H}$ and $u_0^n = \Pi_n u_0$. Recall Example 3.3 and notice that we can apply Theorem 3.1, and by making use of (2.14) we get the estimate

$$E \sup_{0 \leq t \leq T} |u^n(t) - u(t)|_0^2 + E \int_0^T |u^n(t) - u(t)|_1^2 dt \leq C 2^{-2n},$$

with a constant C independent of n . Assume now also **(A5)**, recall Example 4.3 and define $A^{n,\tau}$ and $B^{n,\tau}$ by (4.21). Notice that we can apply Theorem 4.4. Hence if $u_0^{n,\tau} = \Pi_n u(0)$ we get the estimate

$$E \max_{0 \leq i \leq m} |u_i^{n,\tau} - u(i\tau)|_0^2 + \tau E \sum_{0 \leq i \leq m} |u_i^{n,\tau} - u(i\tau)|_1^2 \leq C (\tau + 2^{-2n}). \quad (6.3)$$

Finally recall Example 5.3 and define $A^{n,\tau}$ and $B^{n,\tau}$ as in Example 5.3. Then we can apply Theorem 3.1, and if $u_{\tau,0}^n := \Pi_n u(0)$ and $T2^{2n}/m \leq \gamma$ for some constant $\gamma < c\lambda$, then we get estimate (6.3) for the explicit space-time approximations $u_{\tau,i}^n$, in place of $u_i^{n,\tau}$, with some constant C .

Let us now recall Example 2.7 and approximate (6.1)–(6.2) by finite difference schemes. Consider first the following system of SDEs, corresponding to the space discretization with finite differences for fixed $h \in (0, 1)$:

$$\begin{aligned} dv(t) &= (L_h(t)v(t) + F_h(t, \nabla_h v(t), v(t))) dt \\ &\quad + \sum_k (M_{k,h}(t)v(t) + g_{k,h}(t)) dW^k(t), \quad z \in \mathbb{G} = hZ^d, \end{aligned} \quad (6.4)$$

$$v(0) = (u_0(z))_{z \in \mathbb{G}}, \quad (6.5)$$

where $g_{k,h}(t) = (g_k(t, z))_{z \in \mathbb{G}}$, $F_h(t, p, r) = (F(t, z, p, r))_{z \in \mathbb{G}}$ and

$$L_h(t)\varphi := \sum_{|\alpha| \leq 1, |\beta| \leq 1} \delta_-^\alpha (a^{\alpha\beta}(t, \cdot) \delta_+^\beta \varphi), \quad \nabla_h \varphi := (\delta_1 \varphi, \delta_2 \varphi, \dots, \delta_d \varphi), \quad (6.6)$$

$$M_{k,h}(t)\varphi := \sum_{|\alpha| \leq 1} b_k^\alpha(t) \delta^\alpha \varphi, \quad (6.7)$$

for functions φ defined on \mathbb{G} . It is not difficult to see that taking the triple $V_n := W_{h,2}^1$, $H_n := W_{h,2}^0$, $V_n^* = (W_{h,2}^1)^*$, problem (6.4)–(6.5) can be cast into equation (3.1), and we can easily check that Assumption 3.1 and equation (3.2) hold. Thus (6.4)–(6.5) has a unique continuous $W_{h,2}^0$ -valued solution $v = v^h$ such that for every $h \in (0, 1)$,

$$E \sup_{t \in [0, T]} |v^h(t)|_{h,0}^2 + E \int_0^T |v^h(t)|_{h,1}^2 dt \leq M < \infty.$$

Assume now that $d = 1$. Consider the normal triple $V \hookrightarrow H \equiv H^* \hookrightarrow V^*$ with $V := W_2^1(\mathbb{R})$, $H := W_2^0(\mathbb{R})$ and $V^* \equiv W_2^{-1}(\mathbb{R})$. Notice that Using (2.19) we can see that there is a constant C such that almost surely for all $t \in [0, T]$

$$\begin{aligned} |D^\alpha (a^{\alpha\beta}(t) D^\beta \varphi) - \delta_-^\alpha (a^{\alpha\beta}(t) \delta_+^\beta \varphi)|_{h,0} &\leq Ch |\varphi|_{W_2^3(\mathbb{R})}, \\ |b_k^\alpha(t) D^\alpha \varphi - b_k^\alpha(t) \delta^\alpha \varphi|_{h,0} &\leq Ch |\varphi|_{W_2^2(\mathbb{R})}, \\ |F_h(t, D\varphi, \varphi) - F_h(t, \delta\varphi, \varphi)|_{h,0} &\leq C|h|_{W_2^2(\mathbb{R})} \end{aligned}$$

for all $\varphi \in W_2^3(\mathbb{R})$ and $h \in (0, 1)$. Hence the consistency condition **(Cn)** holds with $\mathcal{V} = W_2^3(\mathbb{R})$ and $\varepsilon_n = h$. Set $\mathcal{H} = W_2^2(\mathbb{R})$. Assume (A1)-(A4) with $l = 2$. Then the assumptions of Theorem 3.1 are satisfied. Thus there is a constant C such that

$$E \sup_{t \in [0, T]} |u(t) - v^h(t)|_{h,0}^2 + E \int_0^T |u(t) - v^h(t)|_{h,1}^2 dt \leq Ch^2$$

for all $h \in (0, 1)$. Now we approximate (6.5) by the following Euler approximation schemes:

$$\begin{aligned} w_{i+1} &= w_i + (L_h(t_{i+1})w_{i+1} + F_h(t_{i+1}, \nabla_h w_{i+1}, w_{i+1})) \tau \\ &\quad + \sum_k (M_{k,h}(t_i)w_i + g_{k,h}(t_i))(W^k(t_{i+1}) - W^k(t_i)), \quad w_0 = u_0, \quad (6.8) \\ u_{i+1} &= u_i + (L_h(t_i)u_i + F_h(t_{i+1}, \nabla_h u_{i+1}, u_i)) \tau \\ &\quad + \sum_k (M_{k,h}(t_i)u_i + g_{k,h}(t_i))(W^k(t_{i+1}) - W^k(t_i)), \quad v_0 = u_0. \end{aligned}$$

for $i = 0, 1, 2, \dots, m-1$, $\tau = T/m$, $t_i = i\tau$. Then by Proposition 4.3 we get the existence of a unique $W_{h,2}^1$ -valued solution w_i of (6.8), such that w_i is \mathcal{F}_{t_i} -measurable for $i = 1, 2, \dots, m$, if τ is sufficiently small. By Theorem 4.4 for $e_i^{h,\tau} = (u(t_i, z) - w_i(z))_{z \in \mathbb{G}}$, we get

$$E \max_{0 \leq i \leq m} |e_i^{h,\tau}|_{W_{h,2}^0}^2 + \tau \sum_{1 \leq i \leq m} E |e_i^{h,\tau}|_{W_{h,2}^1}^2 \leq C(\tau + h^2)$$

with a constant C independent of τ and h . Recall that $\vartheta(n) = \kappa^2/h_n^2$ for any sequence $h_n \in (0, 1)$ by (2.16). Set $e_{i,\tau}^h = (u(t_i, z) - u_i(z))_{z \in \mathbb{G}}$. Then applying Theorem 5.2 we get

$$E \max_{0 \leq i \leq m} |e_{\tau,i}^h|_{W_{h,2}^0}^2 + \tau \sum_{0 \leq i < m} E |e_{\tau,i}^h|_{W_{h,2}^1}^2 \leq Ch^2,$$

with a constant C independent of τ and h , provided (5.3) holds with κ^2/h^2 in place of $\vartheta(n)$. To obtain the corresponding results when $d > 1$ we need more regularity in the space variable from the solution u of (6.1)-(6.2). Assuming more regularity on the data, it is possible to get the required regularity of u . We do not want to prove in this paper further results on regularity of the solutions to (6.1). Instead of that we consider the case of linear equations, i.e., when F does not depend on p and r , since in this case the necessary results on regularity of the solutions are well known in the literature. (See e.g. [9] and [16].)

6.2. Linear stochastic PDEs. We consider again equation (6.1)-(6.2) and assume that $F = F(t, x, p, r)$ does not depend on p and r . We fix an integer $l \geq 0$. Instead of **(A4)** we assume the following.

Assumption (A*4) $F(t, x, p, r) = f(t, x)$ and $g_k(t, x)$ are $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable functions of (t, ω, x) , and their derivatives in x up to order l are $\mathcal{P} \otimes \mathcal{B}(\mathbb{R})$ -measurable functions such that

$$|f(t, \cdot)|_l^2 + \sum_k |g_k(t, \cdot)|_l^2 \leq \eta,$$

where η is a random variable such that $E\eta \leq M$.

Instead of **(A5)** we make the following assumption.

Assumption (A*5) *Almost surely*

$$(i) \sum_k |D^\gamma(b_k^\alpha(t) - b_k^\alpha(s))| \leq K|t - s|^{\frac{1}{2}}, \quad \sum_k |g_k(s) - g_k(t)|_l^2 \leq \eta|t - s|.$$

$$(ii) |D^\gamma(a^{\alpha,\beta}(t) - a^{\alpha,\beta}(s))| \leq K|t - s|^{\frac{1}{2}}, \quad |f(t) - f(s)|_l^2 \leq \eta|t - s|$$

for all $|\gamma| \leq l$, $s, t \in [0, T]$, $x \in \mathbb{R}^d$ and multi-indices $|\alpha| \leq 1$ and $|\beta| \leq 1$, where K is a constant and η is a random variable such that $E\eta \leq M$.

Consider the space-time discretizations with finite differences. The implicit and the explicit approximations, $v^{h,\tau}$ and v_τ^h are given by the systems of equations defined for $i = 0, \dots, m-1$ by

$$\begin{aligned} v^{h,\tau}(t_{i+1}) &= v^{h,\tau}(t_i) + \tau(L_h(t_{i+1})v^{h,\tau}(t_{i+1}) + f(t_{i+1})) \\ &\quad + \sum_k (M_{k,h}v^{h,\tau}(t_i) + g(t_i))(W^k(t_{i+1}) - W^k(t_i)), \end{aligned} \quad (6.9)$$

$$v^{h,\tau}(0, z) = u(0, z), \quad z \in \mathbb{G}, \quad (6.10)$$

and

$$\begin{aligned} v_\tau^h(t_{i+1}) &= v_\tau^h(t_i) + \tau(L_h(t_i)v_\tau^h(t_i) + f(t_i)) \\ &\quad + \sum_k (M_{k,h}v_\tau^h(t_i) + g(t_i))(W^k(t_{i+1}) - W^k(t_i)), \end{aligned} \quad (6.11)$$

$$v_\tau^h(0, z) = u(0, z), \quad z \in \mathbb{G}, \quad (6.12)$$

respectively, where $t_i = i\tau = iT/m$, $v^{h,\tau}(t_i)$ and $v_\tau^h(t_i)$ are functions on \mathbb{G} , $L_h(t)$ and $M_{k,h}(t)$ are defined by (6.6) and (6.7).

Take $H_n := W_{h,2}^0$ and the normal triple $V_n \hookrightarrow H_n \equiv H_n^* \hookrightarrow V_n^*$ with $V_n := W_{h,2}^1$. Then it is easy to see that

$$(L_h(t_i)\varphi, \psi)_n \leq C|\varphi|_{V_n}|\psi|_{V_n}, \quad (M_{k,h}(t_i)\varphi, \psi)_n \leq C|\varphi|_{V_n}|\psi|_{H_n} \quad (6.13)$$

for all $\varphi, \psi \in V_n$, where $(\cdot, \cdot)_n$ denotes the inner product in H_n , and C is a constant depending only on d and the constant K from Assumption **(A3)**. Thus we can define $L_h(t_i)$ and $M_{h,k}(t_i)$ as bounded linear operators from V_n into V_n^* and H_n respectively. Due to (2.17) and (2.18), the restriction of u_0 , $f(t_i)$ and $g_k(t_i)$ onto \mathbb{G} are H_n -valued random variables such that

$$\begin{aligned} E|f(t_i)|_{H_n}^2 &\leq p^2 E|f(t_i)|_l^2, \quad E|g_k(t_i)|_{H_n}^2 \leq p^2 E|g(t_i)|_l^2, \\ E|u_0|_{H_n}^2 &\leq p^2 E|u_0|_l^2, \end{aligned}$$

where p is the constant from (2.17). Moreover,

$$2(L_h(t_i)\varphi, \varphi)_n + \sum_k |M_{h,k}\varphi|_{H_n}^2 \leq -\frac{\lambda}{2}|\varphi|_{V_n}^2 + C|\varphi|_{H_n}^2 \quad (6.14)$$

for all $\varphi \in V_n$, where C is a constant depending only on d and on the constant K from Assumption **(A2)**. Thus using the notation $u_i^{n,\tau} = v^{h,\tau}(t_i)$, $u_{\tau,i}^n = v_\tau^h(t_i)$ and defining

$$A_i^{n,\tau}(\varphi) = L_h(t_i)\varphi + f(t_i), \quad B_{k,i}^{n,\tau}(\varphi) = M_{k,h}(t_i)\varphi + g_k(t_i)$$

for $\varphi \in W_{h,2}^1$, we can cast (6.9)–(6.10) and (6.11)–(6.12) into (4.1) and into (5.2), respectively, and we can see that Assumption 4.1 and condition (4.6) hold. Consequently, by virtue of Proposition 4.3, for sufficiently small τ (6.9)–(6.10) has a unique

solution $\{v^{h,\tau}(t_i)\}_{i=0}^m$, such that $v^{h,\tau}(t_i)$ is a $W_{h,2}^1$ -valued \mathcal{F}_{t_i} -measurable random variable and $E|v_i^{h,\tau}|_{h,2}^2 < \infty$. Furthermore, by virtue of Proposition 5.1, (6.11)–(6.12) has a unique solution $\{v_\tau^h(t_i)\}_{i=0}^m$, such that $v_\tau^h(t_i)$ is a $W_{h,2}^1$ -valued \mathcal{F}_{t_i} -measurable random variable and $E|v_\tau^h(t_i)|_{h,2}^2 < \infty$.

Let $r \geq 0$ be an integer, and assume that

$$l > r + 2 + \frac{d}{2}. \quad (6.15)$$

Then Theorem 4.4 gives the following result.

Theorem 6.1. *Let Assumptions (A1), (A2), (A3), (A*4) and (A*5) hold with l satisfying (6.15). Then for sufficiently small τ*

$$E \max_{1 \leq i \leq m} |v^{h,\tau}(t_i) - u(t_i)|_{h,r}^2 + E \sum_{1 \leq i \leq m} \tau |v^{h,\tau}(t_i) - u(t_i)|_{h,r+1}^2 \leq C(h^2 + \tau)$$

for all $h \in (0, 1)$, where $C = C(r, l, p, \lambda, T, K, M, d, d_1)$ is a constant.

Proof. Take $H_n := W_{h,2}^r$, $H := W_2^{l-2}(\mathbb{R}^d)$, $\mathcal{H} := W_2^l(\mathbb{R}^d)$ and the normal triples

$$V_n \hookrightarrow H_n \equiv H_n^* \hookrightarrow V_n^*, \quad V \hookrightarrow H \equiv H^* \hookrightarrow V^*, \quad \mathcal{V} \hookrightarrow \mathcal{H} \equiv \mathcal{H}^* \hookrightarrow \mathcal{V}^*$$

where $V_n := W_{h,2}^{r+1}$, $V_n^* \equiv W_{h,2}^{r-1}$, $V := W_2^{l-1}(\mathbb{R}^d)$, $V^* \equiv W_2^{l-3}(\mathbb{R}^d)$, $\mathcal{V} := W_2^{l+1}(\mathbb{R}^d)$ and $\mathcal{V}^* \equiv W_2^{l-1}(\mathbb{R}^d) = V$. Then due to (6.15) there is a constant p such that for $\Pi_n := R_h$,

$$|\Pi_n \varphi|_{V_n} \leq p |\varphi|_V,$$

for all $\varphi \in V$, by virtue of (2.20). It is easy to check that (6.13)–(6.14) still hold, and hence (6.9)–(6.10), written as equation (4.1), satisfies Assumption 4.1 and condition (4.6) in the new triple as well. Using (2.20) it is easy to show that due to Assumption (A3)

$$\begin{aligned} |L(t_i)\varphi - L_h(t_i)\varphi|_{V_n^*} &\leq |L(t_i)\varphi - L_h(t_i)\varphi|_{H_n} \leq Ch|\varphi|_{W_2^{l+1}(\mathbb{R}^d)}, \\ \sum_k |M_k(t_i)\varphi - M_{k,h}(t_i)\varphi|_{H_n} &\leq Ch|\varphi|_{W_2^l(\mathbb{R}^d)} \end{aligned}$$

for all $\varphi \in W_2^{l+1}(\mathbb{R}^d)$, where C is a constant depending on d, l, r and on the constant K from Assumption (A3). Hence we can see that (Cn τ) holds with $\varepsilon_n = h$. Due to Assumption (A*5) we have

$$\begin{aligned} |L(t)\varphi - L(s)\varphi|_V^2 &\leq C|t - s|, \quad |f(t) - f(s)|_V^2 \leq \eta|t - s|, \\ \sum_k |M_k(t)\varphi - M_k(s)\varphi|_V^2 &\leq C|t - s|, \quad \sum_k |g_k(t) - g_k(s)|_V^2 \leq \eta|t - s|, \end{aligned}$$

where η is the random variable from Assumption (A*5), and C is a constant depending on d, d_1, l and on the constant K from Assumption (A*5). It is an easy exercise to show that due to Assumptions (A3) and (A*4) condition (R3) from Assumption 2.3 holds. From [9] it is known that under the Assumptions (A1)–(A3) and (A*4) the problem (6.4)–(6.5) has a unique solution u on $[0, T]$, and that u is a continuous $W_2^l(\mathbb{R}^d)$ -valued (\mathcal{F}_t) -adapted stochastic process such that

$$E \sup_{t \in [0, T]} |u(t)|_l^2 + E \int_0^T |u(t)|_{l+1}^2 dt \leq$$

$$CE|u_0|_l^2 + CE \int_0^T \left(|f(t)|_{l-1}^2 + \sum_k |g_k(t)|_l^2 \right) dt,$$

where C is a constant depending on d , d_1 and the constants λ and K from Assumptions **(A1)**, **(A3)** and **(A*4)**. Hence the regularity conditions **(R1)** and **(R2)** in Assumption 2.3 clearly hold. Now we can conclude the proof by applying Theorem 4.4. \square

Let us now investigate the rate of convergence of the explicit space-time approximations. Take the normal triple $V_n \hookrightarrow H_n \equiv H_n^* \hookrightarrow V_n^*$ with $V_n := W_{h,2}^{r+1}$, $H_n := W_{h,2}^r$, and notice that due to Assumption **(A3)**

$$(L(t_i)\varphi, \psi)_n \leq C_1|\varphi|_{V_n}|\psi|_{V_n}, \quad (M_{k,h}(t_i)\varphi, \psi)_n \leq C_{2k}|\varphi|_{V_n}|\psi|_{H_n} \quad (6.16)$$

with some constants C_1 and C_{2k} depending only on d , r and the constant K from Assumption **(A3)**. Set $L_1 = C_1^2$ and $L_2 = \sum_k C_{2k}^2$. Then Theorem 5.2 yields the following theorem, which improves a result from [18].

Theorem 6.2. *Let Assumptions **(A1)**, **(A2)**, **(A3)**, **(A*4)** and **(A*5)** hold with l satisfying (6.15). Let h and τ satisfy*

$$L_1\kappa^2\frac{\tau}{h^2} + 2\kappa(L_1L_2)^{1/2}\frac{\sqrt{\tau}}{h} \leq q \quad (6.17)$$

for a constant $q < \lambda$. Then

$$E \max_{1 \leq i \leq m} |v_\tau^h(t_i) - u(t_i)|_{h,r}^2 + E \sum_{0 \leq i < m} \tau |v_\tau^h(t_i) - u(t_i)|_{h,r+1}^2 \leq C(h^2 + \tau)$$

for all $h \in (0, 1)$, where $C = C(r, l, p, \lambda, q, T, K, M, d, d_1)$ is a constant.

Proof. As in the proof of Theorem 6.1 we take $H_n := W_{h,2}^r$, $H := W_2^{l-2}(\mathbb{R}^d)$, $\mathcal{H} := W_2^l(\mathbb{R}^d)$ and the normal triples

$$V_n \hookrightarrow H_n \equiv H_n^* \hookrightarrow V_n^*, \quad V \hookrightarrow H \equiv H^* \hookrightarrow V^*, \quad \mathcal{V} \hookrightarrow \mathcal{H} \equiv \mathcal{H}^* \hookrightarrow \mathcal{V}^*$$

with $V_n := W_{h,2}^{r+1}$, $V := W_2^{l-1}(\mathbb{R}^d)$, $\mathcal{V} = W_2^{l+1}(\mathbb{R}^d)$, we cast (6.11)–(6.12) into (5.2), and see that Assumptions 2.3 and 4.1, conditions **(Cn τ)** and (4.6) of Theorem 5.2 hold. Furthermore, $\vartheta(n) = \frac{\kappa^2}{h^2}$. We can easily check that by virtue of (6.16) and (2.16), condition (6.17) yields condition (5.3). Hence applying Theorem 5.2 we finish the proof. \square

Corollary 6.3. *Let $k \geq 0$ be an integer and let Assumptions **(A1)**, **(A2)**, **(A3)**, **(A*4)** and **(A*5)** hold with l satisfying $l > k + 2 + d$. Then the following statements are valid for all multi-indices $|\alpha| \leq k$:*

(i) *For sufficiently small τ*

$$E \max_{1 \leq i \leq m} \sup_{z \in \mathbb{G}} |\delta^\alpha(v_\tau^{h,\tau}(t_i, z) - u(t_i, z))| \leq C(h + \sqrt{\tau})$$

holds for all $h \in (0, 1)$, where $C = C(l, p, \lambda, T, K, M, d, d_1)$ is a constant.

(ii) *Assume also that τ and h satisfy (6.17). Then*

$$E \max_{1 \leq i \leq m} \sup_{z \in \mathbb{G}} |\delta^\alpha(v_\tau^h(t_i, z) - u(t_i, z))| \leq C(h + \sqrt{\tau}) \leq C \left(1 + \kappa^{-1} \sqrt{\lambda/L_1} \right) h$$

for all $h \in (0, 1)$, where $C = C(r, l, p, \lambda, q, T, K, M, d, d_1)$ is a constant.

Proof. By the discrete version of Sobolev's theorem on embedding $W_m^2(\mathbb{R}^d)$ into $C^k(\mathbb{R}^d)$ one knows that if $m \geq k + \frac{d}{2}$, then

$$\sup_{z \in \mathbb{G}} |\delta^\alpha \varphi(z)| \leq C |\varphi|_{W_{h,2}^m}$$

for all $h \in (0, 1)$, $\varphi \in W_{h,2}^m$ and $|\alpha| \leq k$, where $C = C(d, m, k)$ is a constant (see e.g. [18]). Hence the above statements follow immediately from Theorems 6.1 and 6.2. \square

Acknowledgments: The authors thank A. Cohen, G. Kerkyacharian and G. Grün for stimulating discussions about wavelets and finite elements. They are also grateful to the referee for his comments and suggestions which helped to improve the presentation of the paper.

REFERENCES

- [1] Ciarlet, P., The finite element method for elliptic problems. Studies in Mathematics and its Applications Vol. 4, North Holland Publishing Company (1978).
- [2] Cohen, A.: Numerical Analysis of wavelet methods. Studies in Mathematics and its Applications 32, Elsevier (2003).
- [3] Gyöngy, I., Krylov, N.V.: On stochastic equations with respect to semi-martingales II, Ito formula in Banach spaces. *Stochastics* 6, 153–173 (1982).
- [4] Gyöngy, I.: On stochastic equations with respect to semimartingales III. *Stochastics* 7, 231–254 (1982).
- [5] Gyöngy, I.: Lattice approximations for stochastic quasi-linear parabolic partial differential equations driven by space-time white noise II. *Potential Analysis* 11, 1–37 (1999).
- [6] Gyöngy, I. Martinez, T.: Solutions of partial differential equations as extremals of convex functionals. *Acta Math. Hungar.*, 127-145 (2005).
- [7] Gyöngy, I., Millet, A.: On Discretization Schemes for Stochastic Evolution Equations. *Potential Analysis* 23, 99–134 (2005).
- [8] Gyöngy, I., Millet, A.: Rate of Convergence of Implicit Approximations for stochastic evolution equations, Stochastic Differential Equations: Theory and Applications (A volume in Honor of Boris L. Rosovskii) Editors P. Baxendale et S. Lototsky, World Scientific Interdisciplinary Mathematical Sciences Volume 2, p. 281-310 (2007).
- [9] Krylov, N.V. Rosovskii, B.L.: On Cauchy problem for linear stochastic partial differential equations. *Math. USSR Izvestija*, Vol. 11 4, 1267-1284 (1977).
- [10] Krylov, N.V. Rosovskii, B.L.: Stochastic evolution equations. *J. Soviet Mathematics*, 16, 1233–1277 (1981).
- [11] Lions, J.L.: Quelques méthodes de résolution des problèmes aux limites non linéaires. Études mathématiques. Dunod Gauthiers-Villars (1969).
- [12] Mueller-Gronbach, T., Ritter, K.: An Implicit Euler Scheme with Non-uniform Time Discretization for Heat Equations with Multiplicative Noise. Available via arXiv math.PR/0604600, April 2006.
- [13] Pardoux, E.: Équations aux dérivées partielles stochastiques nonlinéaires monotones. Étude de solutions fortes de type Itô, Thèse Doct. Sci. Math. Univ. Paris Sud (1975).
- [14] Pardoux, E.: Stochastic partial differential equations and filtering of diffusion processes. *Stochastics* 3-2, 127–167 (1979).
- [15] Pardoux, E.: Filtrage non linéaire et équations aux dérivées partielles stochastiques associées. École d'été de Probabilités de Saint-Flour 1989. Lecture Notes in Math. 1464, Springer Verlag, 67-163 (1981).
- [16] Rozovskii, B.: Stochastic evolution systems. Linear theory and applications to nonlinear filtering. Kluwer, Dordrecht (1990).
- [17] Yan, Y.B.: Semidiscrete Galerkin Approximation for a Linear Stochastic Parabolic Partial Differential Equation Driven by an Additive Noise. *Num. Math.* 44, 829-847 (2004).

- [18] Yoo, H.: An analytic approach to stochastic partial differential equations and its applications. Thesis, University of Minnesota (1998).

Addresses :

Istvan Gyöngy,
School of Mathematics University of Edinburgh,
and Maxwell Institute for Mathematical Sciences,
King's Buildings,
Edinburgh, EH9 3JZ, United Kingdom
email : gyongy@maths.ed.ac.uk

and

Annie Millet
Laboratoire de Probabilités et Modèles Aléatoires
Universités Paris 6-Paris 7, Boîte Courrier 188,
4 place Jussieu, 75252 Paris Cedex 05, France
and SAMOS-MATISSE, Centre d'Économie de la Sorbonne,
Université Paris 1 Panthéon Sorbonne,
90 Rue de Tolbiac,
75634 Paris Cedex 13, France
email : annie.millet@upmc.fr *and* amillet@univ-paris1.fr